



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 




CD79B Recombinant Monoclonal Antibody

Product Code	CSB-RA004958MA2HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P40259
Immunogen	Recombinant Human CD79B protein
Species Reactivity	Human
Tested Applications	ELISA, FC; Recommended dilution: ELISA:1:5000-1:50000, FC:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Affinity-chromatography
Isotype	hIgG1
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Research Area	Immunology
Target Names	CD79B
Clone No.	8E9


Image



The Binding Activity of Human CD79B with Anti-CD79B recombinant antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79A(CSB-MP004958HU1) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC₅₀ is 3.214-3.642 ng/mL.




The Binding Activity of Human CD79A&CD79B with Anti-CD79B recombinant antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79A&CD79B(CSB-MP5861HU) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC₅₀ is 14.85-17.47 ng/mL.




The Binding Activity of Human CD79B with Anti-CD79B recombinant antibody

Activity: Measured by its binding ability in a functional ELISA. Immobilized Human CD79B(CSB-MP004958HU1p7) at 2 µg/mL can bind Anti-CD79B recombinant antibody. The EC_{50} is 31.97-35.69 ng/mL.



Untransfected HEK-293T cells (green line) and transfected Human CD79B HEK-293T stable cells (red line) were stained with anti-CD79B antibody(CSB-RA004958MA2HU) (2µg/1*10⁶cells), washed and then followed by APC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.



Untransfected HEK-293T cells (green line) and transfected Human CD79A&CD79B HEK-293T stable cells (red line) were stained with anti-CD79B antibody(CSB-RA004958MA2HU) (2µg/1*10⁶cells), washed and then followed by APC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.