



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 




PTK7 Recombinant Monoclonal Antibody

Product Code	CSB-RA622651MA1HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q13308
Immunogen	Recombinant Human PTK7 protein
Species Reactivity	Human, Macaca fascicularis
Tested Applications	ELISA; Recommended dilution: ELISA:1:5000-1:50000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Affinity-chromatography
Isotype	hIgG1
Clonality	Monoclonal
Product Type	Recombinant Antibody
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 95% as determined by SEC-HPLC.
Research Area	Tags & Cell Markers; Metabolism; Signal transduction
Target Names	PTK7
Clone No.	8H7


Image



The Binding Activity of Monkey PTK7 with Anti-PTK7 recombinant antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Monkey PTK7(CSB-MP5478MOV) at 2 µg/mL can bind Anti-PTK7 recombinant antibody. The EC₅₀ is 1.399 - 1.666 ng/mL.



The Binding Activity of Human PTK7 with Anti-PTK7 recombinant antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized Human PTK7(CSB-MP622651HU2) at 2 µg/mL can bind Anti-PTK7 recombinant antibody. The EC₅₀ is 0.3675 - 0.6584 ng/mL.



The purity of PTK7 was greater than 95% as determined by SEC-HPLC

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.