



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Common Name	FMC63
Synonyms	CD19;4;VID3;GC12802
Conjugate	Unconjugated
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Mus musculus
IgG type	Mouse IgG2a
Reactivity	Human
Target	CD19
Uniprot ID	P15391
Description	Anti-CD19(FMC63 biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with 1 μ g/mL Anti-CD19 mAb (BME100094) on Expi293 cells transfected with Human CD19 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

Anti-CD19(FMC63 biosimilar) mAb ELISA

0.2 μ g of Human CD19(M75I,L82V,F83L), hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD19 (M75I,L82V,F83L) Protein, hFc Tag (PME101557) can bind Anti-CD19(FMC63 biosimilar) mAb (BME100094) in a linear range of 3.20–400 ng/mL.



Anti-CD19(FMC63 biosimilar) mAb ELISA

0.2 µg of Human CD19(M75V,R76S,F85S), hFc tagged protein per well




Figure 3. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human CD19 (M75V,R76S,F85S) Protein, hFc Tag (PME101558) can bind Anti-CD19(FMC63 biosimilar) mAb (BME100094) in a linear range of 3.20–400 ng/mL.

