



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Common Name	NA
Conjugate	Unconjugated
Synonyms	C1QR1, MXRA4
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Host Species	Humanized
IgG type	IgG1
Reactivity	Human
Target	CD93
Uniprot ID	Q9NPY3
Description	Anti-CD93(biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with 1 μ g/mL Anti-CD93(biosimilar) mAb (BME100176) on Expi293 cells transfected with Human CD93 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD93 Protein, hFc Tag(PME100689) can bind Anti-CD93(biosimilar) mAb(BME100176) in a linear range of 3.20-80 ng/mL. In order to specifically detect BME100176, mouse anti-human Fab-specific antibody was used as detection antibody.






Figure 3. Flow cytometry analysis of antigen binding of anti-human CD93 mAb(BME100176).
(A) BME100176 does not bind to Jurkat cells that do not express CD93.
(B) A clear peak shift of BME100176 was seen compared to the control when incubated with CD93-expressing THP-1 cells, indicating strong binding of BME100176 to CD93. Antibodies were incubated at 5 µg/mL.

