



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	GPR20
Synonyms	G-protein coupled receptor 20
Description	Human GPR20-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q99678
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	N/A
Molecular Weight	The human full length GPR20-Strep protein has a MW of 38.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	GPR20 is one of the orphan GPCRs that has been identified from human genomic DNA by PCR amplification using primers based on the sequences of the opioid/somatostatin-related receptors, GPR7 and GPR8. The expression of human GPR20 has been detected in several brain regions, including the caudate nuclei, putamen, and the thalamus. A recently disclosed patent demonstrated that GPR20-deficient mice exhibited a hyperactivity disorder characterized by an increase in total distance traveled in an open field test, implying a substantial role of GPR20 in neurophysiological function. However, the physiological mechanisms of GPR20 action, including the identification of natural ligands for GPR20, have not yet been elucidated.
Usage	Research use only
Conjugate	Unconjugated

