

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target FXYD7 **Synonyms** N/A

DescriptionHuman FXYD7-Strep full length protein-synthetic

Delivery 6~8weeks
Uniprot ID P58549
Expression Host HEK293

Protein Families Ion Channels: Other

Protein Pathways N/A

Storage & Shipping

Molecular Weight

The human full length FXYD7-Strep protein has a

MW of 8.5 kDa

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Lyophilized from nanodisc solubilization buffer (20

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This reference sequence was derived from multiple replicate ESTs and validated by similar human genomic sequence. This gene encodes a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7 invariant and 6 highly conserved amino acids. The approved human gene nomenclature for the family is FXYD-domain

containing ion transport regulator.

Transmembrane topology has been established for two family members (FXYD1 and FXYD2), with the Naturality outroed living and the Cataminus

the N-terminus extracellular and the C-terminus on the cytoplasmic side of the membrane. FXYD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. This gene product, FXYD7, is novel and has not been characterized as a protein. [RefSeq curation by

Kathleen J. Sweadner, Ph.D.,

sweadner@helix.mgh.harvard.edu., Dec 2000]

Email: info@dimabio.com Website: www.dimabio.com

Usage Research use only

Conjugate Unconjugated



