



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	ACHB4
Synonyms	N/A
Description	Human ACHB4-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P30926
Expression Host	HEK293
Protein Families	Ion Channels: Cys-loop Receptors
Protein Pathways	N/A
Molecular Weight	The human full length ACHB4-Strep protein has a MW of 56.4 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene is found within a conserved gene cluster and encodes one of the beta subunits of the nicotinic acetylcholine receptor (nAChRs) superfamily which form ligand-gated ion channels with a central pore that forms a cation channel. Neuronal nAChRs are pentameric structures that can be either homomeric or heteromeric, with heteromeric structures containing both alpha and beta subunits. Each subunit contains an extracellular amino terminus and four transmembrane domains. Nicotine is one of the agonists that binds to the receptor. Variants in this gene have been associated with nicotine dependence and lung cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2017]
Usage	Research use only
Conjugate	Unconjugated

