



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Target	CCR2
Synonyms	Ckr2; Ccr2a; Ccr2b; Ckr2a; Ckr2b; mJe-r; Cmkbr2; Cc-ckr-2
Description	Recombinant mouse CCR2 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P51683
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	Mouse CCR2(Met1-Ala55) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 32.3 kDa after removal of the signal peptide. The apparent molecular mass of mCCR2-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Enables C-C chemokine binding activity and C-C chemokine receptor activity. Involved in several processes, including leukocyte migration; positive regulation of cell migration; and regulation of cytokine production. Acts upstream of or within several processes, including cellular defense response; monocyte chemotaxis; and neutrophil clearance. Located in external side of plasma membrane. Is expressed in several structures, including alimentary system; brain; genitourinary system; hemolymphoid system gland; and liver and biliary system. Used to study Coronavirus infectious disease and age related macular degeneration. Human ortholog(s) of this gene implicated in several diseases, including Kawasaki disease; aggressive periodontitis; coronary artery disease (multiple); glucose metabolism disease (multiple); and uveitis (multiple). Orthologous to human CCR2 (C-C motif chemokine receptor 2). [provided by Alliance of Genome Resources, Apr 2022]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Mouse CCR2 Protein, hFc Tag on SDS-PAGE under reducing condition.

