



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION

Target	FCRL5
Synonyms	CD307; FCRH5; IRTA2; BXMAS1; CD307e; PRO820
Description	Recombinant human FCRL5(745-850)Protein with N-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q96RD9
Expression Host	HEK293
Tag	N-Human Fc tag
Molecular Characterization	hFc(Glu99-Ala330)FCRL5(Val745-Thr850)
Molecular Weight	The protein has a predicted molecular mass of 37.2 kDa after removal of the signal peptide. The apparent molecular mass of hFc-FCRL5(745-850) is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the immunoglobulin receptor superfamily and the Fc-receptor like family. This gene and several other Fc receptor-like gene members are clustered on the long arm of chromosome 1. The encoded protein is a single-pass type I membrane protein and contains 8 immunoglobulin-like C2-type domains. This gene is implicated in B cell development and lymphomagenesis. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2010]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human FCRL5(745-850) Protein, hFc Tag on SDS-PAGE under reducing condition.

Human FCRL5(745-850), hFc Tagged protein ELISA
0.2 µg of Human FCRL5(745-850), hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human FCRL5(745-850) Protein, hFc Tag (PME101494) can bind Anti-FCRL5(cevostamab biosimilar without anti-CD3 portion) mAb (BME100089) in a linear range of 3.20-400 ng/mL.

