

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Cat. No. PME101508



#### PRODUCT INFORMATION

OSM **Target Synonyms** ONCM

Recombinant human OSM Protein with C-terminal **Description** 

10×His tag

**Delivery** In Stock **Uniprot ID** P13725 **Expression Host HEK293** 

Tag C-10×His tag

Molecular Characterization

OSM(Ala26-Arg221) 10×His tag

The protein has a predicted molecular mass of

23.5 kDa after removal of the signal peptide. The **Molecular Weight** apparent molecular mass of OSM-His is

approximately 25-35 kDa due to glycosylation.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

staining.

Formulation &

**Background** 

Purity

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Lyophilized from sterile PBS, pH 7.4. Normally 5 %

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the leukemia inhibitory factor/oncostatin-M (LIF/OSM) family of

proteins. The encoded preproprotein is

proteolytically processed to generate the mature protein. This protein is a secreted cytokine and growth regulator that inhibits the proliferation of a number of tumor cell lines. This protein also regulates the production of other cytokines,

including interleukin 6, granulocyte-colony stimulating factor and granulocyte-macrophage colony stimulating factor in endothelial cells. This gene and the related gene, leukemia inhibitory factor, also present on chromosome 22, may have resulted from the duplication of a common ancestral gene. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016]

> Email: info@dimabio.com Website: www.dimabio.com

**Usage** Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)









Figure 1. Human OSM Protein, His Tag on SDS-PAGE under reducing condition.



Email: info@dimabio.com Website: www.dimabio.com

