



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Histone H4 (Acetyl Lys12) Monoclonal Antibody

catalog number: **AN301411L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Rat;Mouse
Immunogen	Acetylated human histone H4 (Lys 12) peptide
Host	Rabbit
Isotype	IgG, κ
Clone	A106
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution


WB	1:500-1:1000
IHC	1:50-1:200

Data




Western Blot with Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:1000. (-): MCF-7, (+): MCF-7+ Sodium butyrate (50mM, 24hr) + Trichostatin A (500ng/ml, 4 hr)

Observed-MW:11 kDa
Calculated-MW:11 kDa




Western Blot with Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:1000. (-): NIH/3T3, (+): NIH/3T3+Trichostatin A (500ng/ml,4h)

Observed-MW:11 kDa
Calculated-MW:11 kDa





Western Blot with Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:1000. (-): C6, (+): C6+Trichostatin A (500ng/ml,4h)

Observed-MW:11 kDa
Calculated-MW:11 kDa



Immunohistochemistry of paraffin-embedded Human spleen using Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:200.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1

Immunohistochemistry of paraffin-embedded Mouse pancreas using Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:200.

Immunohistochemistry of paraffin-embedded Rat pancreas using Histone H4 (Acetyl Lys12) Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Histone post-translational modifications (PTMs), known as the “histone code”, are key mechanisms of epigenetics that modulate chromatin structures. The PTMs on histone including acetylation, methylation, phosphorylation, and novel acylations directly affect the accessibility of chromatin to transcription factors and other epigenetic regulators, altering genome stability and gene transcription. Histone acetylation, tightly controlled by the opposing action of histone acetyltransferases (HATs) and histone deacetylases (HDACs), occurs primarily at lysine residues on the N-terminal tails of histones H2A (Lys5, 9, and 15), H2B (Lys5, 12, 15, 16, and 20), H3 (Lys4, 9, 14, 18, 23, 27, and 36), and H4 (Lys5, 8, 12, 16, and 20), and plays vital roles in the regulation of gene expression, DNA damage repair, chromatin dynamics, etc.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.1