



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08135 - Goat Anti-EPB41L2 / 4.1G (aa 593 to 604) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EPB41L2, 4.1-G, erythrocyte membrane protein band 4.1-like 2, RP3-324N14.1, DKFZp781D1972, DKFZp781H1755, OTTHUMP00000040262, erythrocyte membrane protein band 4.1 like-protein 2

Official Symbol: EPB41L2

Accession Number(s): NP_001422.1; NP_001129026.1; NP_001186317.1; NP_001239589.1; NP_001337228.1; NP_001337230.1; NP_001337232.1; NP_001337233.1; NP_001337236.1; NP_001337243.1; NP_001337249.1

Human GeneID(s): [2037](#)

Important Comments: This antibody is reported to recognise all isoforms NP_001422.1 (iso a), NP_001129026.1 (iso b), NP_001186317.1 (iso c), NP_001239589.1 (iso d), NP_001337228.1 (iso e), NP_001337230.1 (iso f), NP_001337232.1 (iso g), NP_001337233.1 (iso h), NP_001337236.1 (iso i), NP_001337243.1 (iso j), NP_001337249.1 (iso k).

Immunogen

Peptide with sequence C-RREVRSPKAPH, from the internal region of the protein sequence according to NP_001422.1; NP_001129026.1; NP_001186317.1; NP_001239589.1; NP_001337228.1; NP_001337230.1; NP_001337232.1; NP_001337233.1; NP_001337236.1; NP_001337243.1; NP_001337249.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Preliminary testing showed bands at approx 110+40kDa in Human Cerebellum lysates after 0.1µg/ml antibody staining (calculated MW of 113kDa according to NP_001422.1). These bands were successfully blocked by incubation with the immunizing peptide. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Uterus and Colon. Recommended concentration: 2.5µg/ml.


Immunofluorescence: Strong expression of the protein seen in the membranes of A431 and in the cell junctions of U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of A431 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human




EB08135 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).




EB08135 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cell junction staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08135 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB08135 (2.5µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB08135 (2.5µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.