



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



CD3 FITC - CD28 PE. Mouse Bi-Test™ Reagents (FITC/RPE)

COMMENTS

The CD3 epitope is expressed on the epsilon chain of the CD3/T cell antigen receptor (TcR) complex. CD3 is expressed on 65-85% of thymocytes and has a mitogenic effect on peripheral blood T cells. Identification of human T cells expressing the 22-28,000 M. W. surface antigen. Antihuman CD28 binds the 44kDa MW cell surface protein on the surface of most T cells. CD28 acts as the ligand for the B7/BB-1 molecule on the surface of activated B cells. B7/BB-1 co-stimulates T cells through CD28, along with CD2 and CD3. CD28 antigen is a disulfide-linked homodimeric glycoprotein. The CD28 antigen is present on approximately 60%-80% of lymphocytes (95% of CD4 and 50% of CD8 lymphocytes). CD28 regulates the expression of cytokines by T cells, not only IL-2, but also IL-1 alpha and CSF-1, usually synthesized by accessory cells. CD28 functions as a cell adhesion molecule (CAM) for certain T cell subsets.

CONCENTRATION

See vial for concentration

SHIP CONDITIONS

Room Temperature

STORAGE CUSTOMER

Product should be stored at 4-8°C. DO NOT FREEZE

STABILITY

Reagents are stable for the period shown on the vial label when stored properly

Use

PBMC: Add 10 µl of MAB/10⁶ PBMC in 100 µl PBS. Mix gently and incubate for 15 minutes at 2° to 8°C. Wash twice with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. **WHOLE BLOOD:** Add 10 µl of MAB/100 µl of whole blood. Mix gently and incubate for 15 minutes at room temperature 20°C. Lyse the whole blood. Wash once with PBS and analyze or fix with 0.5% v/v of paraformaldehyde in PBS and analyze. See instrument manufacturer's instructions for Lysed Whole Blood and Immunofluorescence analysis with a flow cytometer or microscope.

ORDERING INFORMATION

CATALOG NUMBER

0328S

SIZE

50 Tests

FORM

Bi-Test (FITC/RPE) Reagent

HOST/CLONE

Mouse

FORMULATION

Provided as sterile filtered solution in phosphate buffered saline with 0.08% sodium azide and 0.2% carrier protein

ISOTYPE

IgG1 (F)/IgG1 (PE)

APPLICATIONS

Flow Cytometry