



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Neurogenin-1 Blocking Peptide

### BACKGROUND

This peptide is the immunogen used to make Cat # X2758P.

Acts as a transcriptional regulator. Involved in the initiation of neuronal differentiation. Activates transcription by binding to the E box (5'-CANNTG-3'). Associates with chromatin to enhancer regulatory elements in genes encoding key transcriptional regulators of neurogenesis. Efficient DNA binding requires dimerization with another bHLH protein. Expression restricted to the embryonic nervous system.

### ORDERING INFORMATION

**CATALOG NUMBER**  
X2773B

**SIZE**  
50 µg

**CUSTOMER STORAGE**  
Product should be stored at -20°C.  
Aliquot to avoid freeze/thaw cycles

### FORMULATION

Provided as solution in phosphate buffered saline with 0.08% sodium azide

### SHIP CONDITIONS

Ship at ambient temperature, freeze upon arrival

### STABILITY

Products are stable for one year from purchase when stored properly

### COMMENTS

Optimal concentration should be evaluated by serial dilutions.

### INSTRUCTIONS

Use: Incubate antibody neat with at least a 50 fold stoichiometric excess of blocking peptide at 37°C for 20 minutes (molecular weights of peptide and antibody are ~2.5 kDa and ~160 kDa, respectively). Antibody can then be diluted to a concentration suitable for Western blot.

Example: 10 µl or 10 µg of Exalpha's rabbit anti-Neurogenin-1 (Cat. No. X2758P) is added to 10 µg of blocking peptide for a total volume of 20 µl. The mixture is allowed to incubate for 20 minutes at 37°C prior to dilution in suitable buffer (for Western blot, etc.).