



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Murine Anti-Factor VIII

Clone GMA-8025

Factor VIII (FVIII) is a heterodimer consisting of a heavy chain (ranging in mass from 90 to 200 kDa) bound via metal ions to a light chain (80 kDa). In plasma, FVIII circulates in an inactive form bound to von Willebrand factor. Following activation by factor Xa or thrombin, factor VIIIa can function as cofactor for the enzyme factor IXa in the activation of factor X in the presence of phospholipid and Ca²⁺. Absent or defective FVIII is the cause of the X-linked recessive bleeding disorder hemophilia A. GMA-8025 recognizes the C2 domain of FVIII, and is suitable for ELISA and Western blot applications.

Description

Antibody Source:	mouse monoclonal, IgG _{2a}
Antigen Species Bound:	human
Specificity:	FVIII C2 domain
Immunogen:	B-domain deleted recombinant human FVIII

Formulation and Storage

Purity:	Purified by protein G affinity chromatography from serum-free cell culture supernatant.
Product Formulation:	Lyophilized from a ≥1 mg/ml solution in 20 mM NaH ₂ PO ₄ 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 (ε _{0.1%}).
Reconstitution:	Reconstitute with deionized water.
Storage:	Store lyophilized or reconstituted and aliquoted material at -20° C for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at 4° C.
Country of Origin:	USA
Size Options:	0.1 mg or 0.5 mg

Applications

Working Concentration:	Approximately 1-5 µg/ml. Researcher should titer antibody in specific assay.
ELISA:	Binds immobilized human FVIII.
Immunoblotting:	Western blot detects C2 domain of human FVIII under reduced conditions.

