



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CD38 antibody [MSVA-038R] HistoMAX™

Cat. No. GTX04967

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Application	IHC-P
Reactivity	Human

Package

500 µl, 100 µl

PRODUCT

Summary This antibody was validated on 76 different Normal Tissues by IHC-P.

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
--------------------	----------------------

IHC-P	1:50 - 1:100
-------	--------------

Note : Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Tris-EDTA-based Target Retrieval Solution buffer.

Not tested in other applications.

PROPERTIES

Form	Liquid
Buffer	10mM PBS, BSA (Please contact us for sodium azide-free or sodium azide and BSA-free format)
Preservative	sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 µg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide corresponding to residues within 190-310 aa of human CD38
Purification	Protein A/G purified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX04967 IHC-P Image

IHC-P analysis of human appendix tissue using GTX04967 CD38 antibody [MSVA-038R] HistoMAX™. In the appendix CD38 staining occurs in various inflammatory cells while the epithelial cells remain CD38 negative.



GTX04967 IHC-P Image

IHC-P analysis of human prostate tissue using GTX04967 CD38 antibody [MSVA-038R] HistoMAX™. Strong membranous CD38 staining of prostatic epithelial cells.



GTX04967 IHC-P Image

IHC-P analysis of human prostatic adenocarcinoma (PAC) tissue (Gleason 3+3=6) using GTX04967 CD38 antibody [MSVA-038R] HistoMAX™. CD38 negative prostatic adenocarcinoma (Gleason 3+3=6) adjacent to CD38 positive normal epithelium.



For full product information, images and publications, please visit our [website](https://www.genetex.com).