



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MUC2 antibody [C3], C-term

Cat No. GTX100664

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	ICC/IF, IHC-P, IHC-Fr, IHC
Reactivity	Human, Mouse

Reference (8)

Package

25 µl, 100 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IHC-Fr	Assay dependent
IHC	Assay dependent

Not tested in other applications.


Calculated MW 540 kDa. ([Note](#))

PROPERTIES

Form	Liquid
Buffer	1XPBS, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative.
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.48 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human MUC2. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES

GTX100664 IHC-P Image

MUC2 antibody [C3], C-term detects secreted MUC2 protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse colon.

MUC2 stained by MUC2 antibody [C3], C-term (GTX100664) diluted at 1:2000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX100664 IHC-P Image

MUC2 antibody [C3], C-term detects secreted MUC2 protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse intestine.

MUC2 stained by MUC2 antibody [C3], C-term (GTX100664) diluted at 1:2000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTX100664 IHC-P Image

MUC2 antibody [C3], C-term detects secreted MUC2 protein in human intestine by immunohistochemical analysis.

Sample: Paraffin-embedded human intestine.

MUC2 antibody [C3], C-term (GTX100664) diluted at 1:750.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](https://www.genetex.com).