



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Flotillin 1 antibody [HL4250]

Cat. No. GTX643486

Host	Rabbit
Clonality	Monoclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
IHC-P	1:100-1:1000

Not tested in other applications.

Observed MW (kDa) 47 kDa.

Product Note Highly recommended for IHC-P in mouse tissues.

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml
Immunogen	Synthetic peptide of human Flotillin 1.
Purification	Affinity purified by Protein A.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTx643486 IHC-P Image


Flotillin 1 antibody [HL4250] detects Flotillin 1 protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.

Flotillin 1 stained by Flotillin 1 antibody [HL4250] (GTx643486) diluted at 1:200.

Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.



GTx643486 IHC-P Image

Flotillin 1 antibody [HL4250] detects Flotillin 1 protein by immunohistochemical analysis.


Sample: Paraffin-embedded mouse heart.

Flotillin 1 stained by Flotillin 1 antibody [HL4250] (GTx643486) diluted at 1:200.

Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min


GTx643486 WB Image

Various whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with Flotillin 1 antibody [HL4250] (GTx643486) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.


GTx643486 IHC-P Image


Flotillin 1 antibody [HL4250] detects Flotillin 1 protein by immunohistochemical analysis.

Sample: Paraffin-embedded rat tissues.

Flotillin 1 stained by Flotillin 1 antibody [HL4250] (GTx643486) diluted at 1:200.

Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min

Corresponding RNA levels (RPKM) in the tissues are based on NCBI database.


GTx643486 IHC-P Image

Flotillin 1 antibody [HL4250] detects Flotillin 1 protein by immunohistochemical analysis.

Sample: Paraffin-embedded rat heart.

Flotillin 1 stained by Flotillin 1 antibody [HL4250] (GTx643486) diluted at 1:200.

Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min



For full product information, images and publications, please visit our [website](https://www.genetex.com).