



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Nanog antibody, Internal

Cat No. GTX89490

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, ELISA, IHC
Reactivity	Human, Pig

Package
100 µg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
WB	0.03-0.1µg/ml
ICC/IF	Assay dependent
ELISA	Assay dependent
IHC	Assay dependent

Not tested in other applications.

Calculated MW 35 kDa. ([Note](#))

PROPERTIES


Form	Liquid
Buffer	Tris saline pH7.3, 0.02% sodium azide, 0.5% BSA.
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-QNQRMKSKRWQKNN, from the internal region of the protein sequence according to NP_079141.2.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Date 2019 / 07 / 08 Page 1 of 2

DATA IMAGES




GTX89490 WB Image

WB analysis of human ovary lysate using GTX89490 Nanog antibody, Internal.

Dilution : 0.03µg/ml

Loading : 35µg protein in RIPA buffer



GTX89490 ICC/IF Image

ICC/IF analysis of induced pluriform stem cells derived from human keratinocytes using GTX89490 Nanog antibody, Internal.

Green : Primary antibody

Dilution : 5µg/ml



For full product information, images and publications, please visit our [website](#).