



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Clik1 / STK35 antibody, C-term

Cat No. GTX89642

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P, FACS
Reactivity	Human, Mouse, Pig

Package
100 µg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Dilution
WB	1-3µg/ml
ICC/IF	10µg/ml
IHC-P	10µg/ml
FACS	Assay dependent

Note : Strong expression of the protein seen in the nucleus of U2OS cells.

Human Testis shows granular cytoplasmic staining of Leydig cells.

Not tested in other applications.

Calculated MW 58 kDa. ([Note](#))


PROPERTIES

Form	Liquid
Buffer	Tris saline pH7.3, 0.02% sodium azide, 0.5% BSA.
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence ELETRMDQVTCAA, from the C Terminus of the protein sequence according to NP_543026.2.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Date 2019 / 07 / 08 Page 1 of 2

DATA IMAGES

GTx89642 FACS Image


FACS analysis of PFA fixed HEK293 cells using GTx89642 Clik1 / STK35 antibody, C-term.

Blue : Primary antibody

Black : Isotype control

Permeabilization : 0.5% Triton


Dilution : 10µg/ml


GTx89642 WB Image

WB analysis of mouse (A) and pig (B) testes lysate using GTx89642 Clik1 / STK35 antibody, C-term.

Dilution : 1µg/ml


Loading : 35µg protein in RIPA buffer


GTx89642 WB Image

WB analysis of HEK293 nuclear (A) and human breast cancer (B) lysate using GTx89642 Clik1 / STK35 antibody, C-term.

Dilution : 2µg/ml

Loading : 35µg protein in RIPA buffer


GTx89642 ICC/IF Image

ICC/IF analysis of PFA fixed U2OS cells using GTx89642 Clik1 / STK35 antibody, C-term.

Green : Primary antibody

Blue : DAPI

Negative control : Unimmunized goat IgG

Permeabilization : 0.15% Triton

Dilution : 10µg/ml



For full product information, images and publications, please visit our [website](http://www.genetex.com).