

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Anti-Mouse CD120a In Vivo Antibody – Low Endotoxin (55R-593) [ICH1069]

Description

Bulk anti-CD120a In Vivo Antibody – Low Endotoxin (55R-593)

Product Benefits:

Target:

ichorbio's anti-CD120a In Vivo Antibody – Low Endotoxin (55R-593) is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from our <u>competitors</u> at less than 1.0 EU/mg. If ichorbio's low endotoxin antibodies are not low enough we also offer ultra low endotoxin antibodies which have even less endotoxin (<0.75EU/mg) at an even higher purity (98% versus 95%). ichorbio: the best antibodies for *in vivo* research.

CD120a
Clone:
55R-593
Isotype:
Armenian Hamster IgG1
Other Names:
Tumor necrosis factor receptor superfamily member 1A, Tnfrsf1a, TNF-R1, TNF-RI, p55, p60
Uniprot:



P25118

Host:
Armenian Hamster
Species Reactivity:
Mouse
Specificity:
Anti-CD120a In Vivo Antibody – Low Endotoxin (55R-593) recognizes Mouse CD120A
Purification Method:
This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.
Antigen Distribution:
Variety of cell types at low levels
Background:
Tumor Necrosis Factor Receptor Type I (TNFRI) is a 55kD Type I transmembrane protein. TNFRI is expressed independently of TNFRII at low levels on a wide variety of cell types. TNFRI binds TNF-A and TNF-B (also known as LT-A). The subsequent signaling is known to be important for inducing cytolytic activity, antiviral activity, expression of manganous superoxide dismutase and intercellular adhesion molecule (ICAM), IL-6 mRNA accumulation and NF-?B induction. TNFRI carries an approximately 80 amino acid death domain near its carboxy terminus capable of transmitting an apoptotic signal through its interaction with TRADD (TNF Receptor I associated death domain protein). Binding of 55R-170 antibody to TNFRI has been shown to block in vitro and in vivo receptor signaling initiated by ligand binding.
Immunogen:
E. coli-expressed extracellular domain of the mouse TNFRI protein
Concentration:
1.0 mg/ml
Formulation:
0.01 M phosphate buffered saline (PBS) pH 7.2, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and Azide free.



>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

Endotoxin:

? 1.0 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

Aggregation:

Aggregation level ? 5%

Aggregation level ? 1%

Storage:

This antibody is stable for at least 4 weeks when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at – 20°C or -80°C. Avoid repeated freeze thaw cycles.

Applications:

Immunoprecipitation, Western Blot

Application Notes:

Each investigator should determine their own optimal working dilution for specific applications.

Use:

Products are for research use only. Not for use in diagnostic or therapeutic procedures.

Isotype Control:

Armenian Hamster IgG Isotype Control for In Vivo – Low Endotoxin [PIP] (ICH2251)

Antibodies against the same target:

Anti-CD120a In Vivo Antibody – Low Endotoxin [55R-170] (ICH1094)