



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Anti-Mouse CD105 (MJ7/18) In Vivo Antibody- Low Endotoxin

Anti-Mouse CD105 In Vivo Antibody - Low Endotoxin (MJ7/18)

Product Benefits:

ichorbio's anti-Mouse CD105 In Vivo Antibody (MJ7/18) - Low Endotoxin [ICH1191] is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from [Bio X Cell](#) at less than 1.0 EU/mg. If ichorbio's low endotoxin antibodies are not low enough we also offer ultra low endotoxin antibodies which have even less endotoxin (0.5EU/mg) at an even higher purity (98% versus 95%). ichorbio: the best antibodies for *in vivo* research.

Target:

CD105

Clone:

MJ7/18

Size:

ichorbio's MJ7/18 *in vivo* antibody is available in the following bulk sizes: 5mg, 25mg, 50mg and 100mg. ichorbio regularly manufactures multi-gram amounts of our anti-mouse CD105 MJ7/18 clone - please contact us for pricing.

Isotype:

IgG2a kappa

Other Names:

Endoglin, Ancillary TGF-beta Receptor

Host:

Rat

Species Reactivity:

Mouse

Specificity:

MJ7/18 activity is directed against mouse CD105 (endoglin).

Antigen Distribution:

CD105 is alternatively spliced, resulting in both long (L-endoglin) and short (S-endoglin) forms. The predominant isoform, L-endoglin, promotes the proliferation and migration of endothelial cells via enhanced ALK1-Smad1/5 signaling and is the presumed isoform when not specified in the literature. In contrast, S-endoglin enhances ALK5-Smad2/3 signaling. Endoglin has specific receptor-ligand interactions between type I and type II TGF- β superfamily receptors when binding TGF- β superfamily ligands. Mouse CD105 has three separate regions of similarity to TGF- β receptor III but does not contain the RGD tripeptide found in human CD105.

Background:

CD105 (endoglin) is a TGF- β superfamily co-receptor that promotes angiogenesis, is involved in endothelial integrin-mediated mural cell and leukocyte adhesion, antagonizes TGF- β mediated ERK activation, is essential to the immune response of macrophages, regulates trophoblast differentiation and invasion during pregnancy, promotes T-cell proliferation, and regulates differentiation and collagen expression in myofibroblasts.

Immunogen:

MJ7/18 was produced by immunizing rats with inflamed mouse skin and selecting for reactivity with endothelial cells. MJ7/18 predominantly stains vascular endothelial cells and is a marker of mouse endothelium

Concentration:

1.0 - 5.0 mg/ml

Formulation:

0.01 M phosphate buffered saline (PBS) pH 7.2, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and Azide free.

Purity:

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

Endotoxin:

1.0 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

Aggregation:

Aggregation level ? 5%

Aggregation level ? 1%

IMPACT Pathogen Test:

We use the IMPACT test generated by IDEXX Laboratories to guarantee our Ultra Low Endotoxin antibodies are pathogen free. Our mouse antibodies are tested for: Mycoplasma spp., Mycoplasma pulmonis, Sendai virus, Mouse

hepatitis virus, Pneumonia virus of mice, Minute virus of mice, Mouse parvovirus (MPV1-5), Theiler's murine encephalomyelitis virus, Murine norovirus, Reovirus 3, Mouse rotavirus, Ectromelia virus, Lymphocytic choriomeningitis virus, Polyoma virus, Lactate dehydrogenase-elevating virus, Mouse adenovirus (MAD1, MAD2), Mouse cytomegalovirus K virus, Mouse thymic virus, Hantaan virus, Corynebacterium bovis, Corynebacterium spp. (HAC2)

Storage:

Anti-Mouse CD105 In Vivo Antibody (MJ7/18) is stable for at least one week when stored sterile at 2-8°C. For long term storage aseptically aliquot in working volumes without diluting and store at -20°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

Applications:

Agonist, Cell Separation, ELISA, Flow Cytometry, Immunofluorescence, IF Microscopy, Immunohistochemistry, IHC FF, Immunoprecipitation, LCI, Western Blot

Application Notes: Each investigator should determine their own optimal working dilution for specific applications. **Use:**

Products are for research use only.

Isotype Control:

[Rat IgG2a In Vivo isotype control - Low Endotoxin \[ICH2244\]](#)

Alternative Names:

- AI528660 antibody
- AI662476 antibody
- CD 105 antibody
- CD105 antibody
- CD105 antigen antibody
- EGLN_HUMAN antibody
- END antibody
- Endoglin antibody
- Eng antibody
- FLJ41744 antibody
- HHT1 antibody
- ORW antibody
- ORW1 antibody
- Osler Rendu Weber syndrome 1 antibody
- RP11 228B15.2 antibody
- S endoglin antibody
- S-endoglin antibody
- SN6 antibody