



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Anti-Human CD147 (VJ1/9)



FITC

147FI-100T

100 test

PE

147PEI-100T

100 test



## 1. PRODUCT DESCRIPTION

- **Clone:** VJ1/9;
- **Isotype:** IgG1;
- **Tested application:** flow cytometry;
- **Immunogen:** The anti-CD147 monoclonal antibody derives from activated HUVEC cells;
- **Species reactivity:** Human;
- **Storage instruction:** store in the dark at 2-8 °C;
- **Storage buffer:** aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN<sub>3</sub>);
- **Recommended usage:** Immunostep's CD147, clone VJ1/9, is a monoclonal antibody intended for the identification and enumeration of leukocytes, erythrocytes, platelets, and endothelial cells using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using 1 test for 10<sup>6</sup> cells;
- **Presentation:** liquid;
- **Source:** Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
- **Purification:** Affinity chromatography;
- **Other names:** Neurothelin, Basigin, EMMPRIN;
- **Gene ID:** 682;
- **Molecular weight:** 50 - 65 kDa.

## 2. ANTIGEN DETAILS

**Large description:** This antibody reacts with the CD147 antigen which reacts with Basigin or Neurothelin, a transmembrane glycoprotein (50-60kd) of the immunoglobulin super-gene family. Neurothelin is a molecule which is broadly expressed on cells of hematopoietic and non-hematopoietic origin but primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. Neurothelin is a blood brain barrier-specific molecule. Its expression on specific cell types may be regulated by cytokines. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.<sup>(1-5)</sup>

## 3. WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

## 4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to [www.immunostep.com](http://www.immunostep.com) technical support for more information.

## 5. REFERENCES

1. Gao J, Hu Z, Liu J, Liu D, Wang Y, Cai M, et al. Expression of CD147 and Lewis y antigen in ovarian cancer and their relationship to drug resistance. *Med Oncol* May;31(5):920.
2. Sudou A, Ozawa M, Muramatsu T. Lewis X structure increases cell substratum adhesion in L cells. *J Biochem*1995 Feb;117(2):271-5.
3. Fadool JM, Linsler PJ. Evidence for the formation of multimeric forms of the 5A11/HT7 antigen. *Biochem Biophys Res Commun*1996 Dec 04;229(1):280-6.
4. Ikeda E, Flamme I, Risau W. Developing brain cells produce factors capable of inducing the HT7 antigen, a blood-brain barrier-specific molecule, in chick endothelial cells. *Neurosci Lett*1996 May 17;209(3):149-52.
5. Schuster VL, Lu R, Kanai N, Bao Y, Rosenberg S, Prie D, et al. Cloning of the rabbit homologue of mouse 'basigin' and rat 'OX-47': kidney cell type-specific expression, and regulation in collecting duct cells. *Biochim Biophys Acta*1996 Mar 27;1311(1):13-9.

## 6. EXPLANATION OF SYMBOLS



Fluorochrome



Product reference



Content for <n> analysis



Regulatory Status



Research Use Only



Manufacturer

## 7. MANUFACTURED BY: IMMUNOSTEP S.L.



**Address:** Avda. Universidad de Coimbra, s/n  
Cancer Research Center (C.I.C)  
Campus de Unamuno  
37007 Salamanca (Spain)  
(+34) 923 294 827  
**Tel./fax:**  
**E-mail:** [info@immunostep.com](mailto:info@immunostep.com)  
[www.immunostep.com](http://www.immunostep.com)