

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Anti-Human CD35 (E11)















35F2-100T

100 test



PRODUCT DESCRIPTION

- Clone: E11;
- Isotype: IgG1;
- Tested application: flow cytometry;
- Immunogen: The anti-CD35 monoclonal antibody derives from human culture cells;
- Species reactivity: Human;
- Storage instruction: store in the dark at 2-8 °C;
- Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₂);
- Recommended usage: Immunostep's CD35, clone E11, is a monoclonal antibody intended for the identification and enumeration of granulocytes, monocytes, B cells, erythrocytes, follicular dendritic cells, subsets of NK and T cells using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for 106 cells;
- Presentation: liquid:
- Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
- Purification: Affinity chromatography;
- Other names: Complement receptor type 1, C3b/C4b receptor, CR1;
- Molecular weight: 223,63 kDa.

2. ANTIGEN DETAILS

Large description: CD35 is a type I single chain of glycoprotein, also known as C3b/C4b receptor. Complement Receptor type 1 or CRI. Four molecular weight allotypes (160kD, 190kD, 220kD, and 250kD) have been described. CD35 is expressed on granulocytes, monocytes, B cells, erythrocytes. follicular dendritic cells, as well as the subsets of NK and T cells, CD35 binds complement C3b. C4b, or iC3, and iC4, and plays important roles in both innate and adoptive immune response via mediating phagocytosis by granulocytes and monocytes. CD35 has also been reported to inhibit T-cell proliferation.(1, 2)

WARRANTY 3.

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, therapeutic or commercial purposes.

Please, refer to www.immunostep.com technical support for more information.

5. REFERENCES

- Dougherty GJ, Selvendran Y, Murdoch S, Palmer DG, Hogg N, The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, IO.I, Eur J Immunol1987 Oct;17(10):1453-9.
- Schlossman SF. Leucocyte typing V: white cell differentiation antigens: proceedings of the Fifth International Workshop and Conference: held in Boston, USA, 3-7 November, 1993. Oxford: Oxford University Press: 1995.

6. **EXPLANATION OF SYMBOLS**

\ .	Fluorochrome
REF	Product reference
Σ	Content for <n> analysis</n>
	Regulatory Status
RUO	Research Use Only
•••	Manufacturer

MANUFACTURED BY:

7.

IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

Campus de Unamuno 37007 Salamanca (Spain)

Telf./fax: (+34) 923 294 827 E-mail: info@immunostep.com

www.immunostep.com