

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Anti-Human CD44 (HP2/9)



1. PRODUCT DESCRIPTION

- Clone: HP2/9;
- Isotype: IqGl;
- Tested application: flow cytometry;
- Immunogen: T cell leukemia HPB-ALL;
- Species reactivity: Human;
- Storage instruction: store in the dark at 2-8 °C;
- Storage buffer: aqueous buffered solution containing protein stabilizer and 0.09% sodium azide (NaN₂);
- Recommended usage: Immunostep's CD44, clone HP2/9 is a monoclonal antibody intended for the identification and enumeration of leucocytes and erythrocytes using flow cytometry. This reagent is effective for direct immunofluorescence staining of human tissue for flow cytometric analysis using I test for I0⁶ cells;
- Presentation: liquid;
- Source: Supernatant proceeding from an in vitro cell culture of a cell hybridoma;
- Specificity: The antibody specifically recognizes the human protein CD44 and it was
 validated by the 5th International Workshop on Human Leukocyte Differentiation
 Antigens (Boston, USA, 3-7 November 1993). Clone HP2/9 is capable of blocking the
 binding of hyaluronic acid to cells^(IA) in its standard isoform (CD44s), probably binding to
 more than one site (5) on CD44.
 - Clone HP2/9 is capable of blocking clone L178 but not clone HI44a, thereby recognizing different epitopes than clone HI44a. L178 clone does not block HP2/9 clone.
- Purification: Affinity chromatography:
- Other names: CDw44, Epican, Extracellular matrix receptor III, ECMR-III, GP90 lymphocyte homing/adhesion receptor, HUTCH-I, Heparan sulfate proteoglycan, Hermes antigen, Hyaluronate receptor, Phagocytic glycoprotein I, PGP-I, Phagocytic glycoprotein I, Pgp-I, H-CAM, qp85, Ly-24;
- Gene ID: 960;
- Molecular weight: 80 95 kDa.

2. ANTIGEN DETAILS

Large description: The monoclonal antibody is directed against the CD44- antigen, which is mainly expressed on leucocytes and erythrocytes. Very few types of tissue or cells lack CD44. Cells and tissues that lack CD44 include platelets, hepatocytes, certain lymphoid cell lines, cardiac muscle, kidney tubular epithelium, testis and portions of the skin. The CD44 antigen is present on approximately 90% of lymphocytes, monocytes, and granulocytes and in lower amounts on thymocytes, fibroblasts, and erythrocytes.

The antigen mediates a variety of functions, including leucocyte–endothelial cell binding and lymphocyte homing to certain peripheral lymphoid microenvironments.⁽ⁱ⁻³⁾

WARRANTY

Warranted only to conform to the quantity and contents stated on the label or in the product labelling at the time of delivery to the customer. Immunostep disclaims hereby other warranties.

Immunostep's sole liability is limited to either the replacement of the products or refund of the purchase price.

4. ADDITIONAL INFORMATION

For research use only. Not for diagnostic use.

Not for resale. Immunostep will not be responsible of violations that may occur with the use of this product. Any use of this product other than the specified in this document is strictly prohibited.

Unless otherwise indicated by Immunostep by written authorization, this product is intended for research only and is not to be used for any other purpose, including without limitation, for human or animal diagnostic, the

Please, refer to www.immunostep.com technical support for more information.

REFERENCES

- Telen MJ, Udani M, Washington MK, Levesque MC, Lloyd E, Rao N. A blood group-related polymorphism of CD44 abolishes a hyaluronan-binding consensus sequence without preventing hyaluronan binding. J Biol Chem. 1996 Mar 22;271(12):7147-53. doi: 10.1074/ ibc.271.12.7147. PMID: 8636151.
- de la Hera A, Acevedo A, Marston W, Sanchez-Madrid F. Function of CD44(Pgp-I) homing receptor in human T cell precursors. Int Immunol. 1989;1(6):598-604. doi: 10.1093/ intimm/1.6.598. PMID: 2489047.
- Lesley J, Hyman R, Kincade PW. CD44 and its interaction with extracellular matrix. Adv Immunol1993:54:271-335.
- García-Peydró M, Fuentes P, Mosquera M, García-León MJ, Alcain J, Rodríguez A, García de Miguel P, Menéndez P, Weijer K, Spits H, Scadden DT, Cuesta-Mateos C, Muñoz-Calleja C, Sánchez-Madrid F, Toribio ML. The NOTCHI/CD44 axis drives pathogenesis in a T cell acute lymphoblastic leukemia model. J Clin Invest. 2018 Jul 2;128(7):2802-2818. doi: 10.1172/JC192981. Epub 2018 May 21. PMID: 29781813; PMCID: PMC6025994.
- Liao HX, Lee DM, Levesque MC, Haynes BF. N-terminal and central regions of the human CD44 extracellular domain participate in cell surface hyaluronan binding. J Immunol. 1995 Oct 15;155(8):3938-45. PMID: 7561101.
- Ugarte-Berzal E, Bailón E, Amigo-Jiménez I, Albar JP, García-Marco JA, García-Pardo A. A novel CD44-binding peptide from the pro-matrix metalloproteinase-9 hemopexin domain impairs adhesion and migration of chronic lymphocytic leukemia (CLL) cells. J Biol Chem. 2014 May 30;289(22):15340-9. doi: 10.1074/jbc.M114.559187. Epub 2014 Apr 16. PMID: 24739387; PMCID: PMC4140891.

6. EXPLANATION OF SYMBOLS

	Fluorochrome
REF	Product reference
\sum	Content for <n> analysis</n>
	Regulatory Status
RUO	Research Use Only
•••	Manufacturer

7. MANUFACTURED BY:

IMMUNOSTEP S.L.

Address: Avda. Universidad de Coimbra, s/n Cancer Research Center (C.I.C)

Campus de Unamuno 37007 Salamanca (Spain)

immunostep

Telf./fax: (+34) 923 294 827
E-mail: info@immunostep.com

www.immunostep.com