

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Lifespan Technologies PO Box 3352 Salt Lake City, UT 84110 Tel: 801-971-1818 Fax: 801-464-6116 info@lifespantech.com

Dextran Sulfate (5000 or 500,000 MW) ELISA Kit for Buffer Samples

INTENDED USE: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT INTENDED FOR CLINICAL OR DIAGNOSTIC USE.

Kit includes: Coated 96-well plate Detector -Enzyme Conjugate Solution TMB Solution Stop Solution Wash Concentrate 10X, (dilute 1 part plus 9 parts)


Researcher must provide:

Pipettes (8 Channel Multipipettor is recommended) Absorbance microplate reader Dextran sulfate standards Tris Buffered Saline (TBS) pH 7.5 (10mM Tris 150mM NaCl) Plate Cover

Storage and Stability

Kit can be stored unopened at 4°C for up to six months. The Detector-Enzyme Conjugate Solution and the TMB solution should be protected from light.

Background



Dextran Sulfate is in the family glycosaminoglycan. It is a polyanionic dextran derivative which may be synthesized from various high purity and well-characterized dextran fractions. In clinical research, anticoagulant dextran sulfate properties have been tested as a possible substitute for heparin in anticoagulant therapy. Another source of interest relates to the effect of dextran sulfate on enzyme inhibition in certain biological systems. Dextran sulfate is used to precipitate LDL and VLDL in plasma fractionation procedures. Dextran sulfate must then be removed from the product. The S-3900 assay allows measurement of extremely low levels of dextran sulfate and gives manufacturers

quantitative data that they have removed dextran sulfate from their product.


The dextran sulfate ELISA product number S-3900 is a quantitative enzyme-linked assay designed for the *in vitro* measurement of dextran sulfate levels in low protein content fluids such as buffer or urine. This assay measures dextran sulfate directly using a dextran sulfate binding protein which has been conjugated to HRP.

The dextran sulfate-ELISA is a competitive assay in which the colorimetric signal is inversely proportional to the amount of dextran sulfate present in the sample. Samples to be assayed are first mixed with the detector-enzyme conjugate in wells of the coated plate. Dextran sulfate in the sample competes with dextran sulfate bound to the plate for binding of the detector-enzyme conjugate. The concentration of dextran sulfate in the sample is determined using a standard curve of known amounts of dextran sulfate.

Reagent Preparation

<u>Dextran Sulfate Standards</u>: Make dilutions of your dextran sulfate using Tris Buffered Saline (TBS) pH 7.5 (10mM Tris 150mM NaCl) to obtain standards of 0.003, 0.01, 0.03, 0.1, 0.3, and 1.0 μ g/mL. Standardization should be performed using dextran sulfate that is the same dextran sulfate type contained in your unknowns.

1X Wash Buffer: Make a 1:10 dilution of 10X Wash Buffer in distilled or deionized water.



Assay Procedure

Dextran Sulfate ELISA

- Set up the dextran sulfate ELISA plate as illustrated above. We suggest the dextran sulfate standard dilution series be run in triplicate for best results. Add 50 μL of Standards and samples into corresponding wells. Add 50 μL of Detector -Enzyme Conjugate to all wells except the Blank wells. Mix well. Cover plate and incubate for 30 minutes at room temperature. A rotator is highly recommended if available.
- Discard the solution and wash the wells four times with 300 μL per well of 1X Wash Buffer. An automated plate washer is
 recommended if available. After washing, immediately proceed to the next step. Do not delay in removing wash buffer from
 the wells. Do not allow plate to dry.
- Add 100 μL TMB Solution to each well. Incubate the plate in the dark at room temperature for 4-60 minutes waiting for the zero dextran sulfate wells to develop to a medium to dark blue color. Watch for color development and DO NOT overdevelop.
- 4. Add 50 µL Stop Solution which will change the color from blue to yellow.
- 5. Immediately measure the absorbance of each well at 450 nm.
- 6. Calculate the binding percentage for each sample using the formula:

 $[A_{450}(Sample) - A_{450}(Blank)] / [A_{450}(Zero \ dextran \ sulfate) - A_{450}(Blank)] x \ 100 = \% \ Binding$

Using linear or nonlinear regression, plot a standard curve of percent binding versus concentration of dextran sulfate standards. Determine dextran sulfate levels of unknowns by comparing their percentage of binding relative to the standard curve. Dextran sulfate can be estimated by comparing the values from the wells containing unknowns to the values in the standard curve.

Lifespan Technologies products are sold for research and development purposes only and are not for diagnostic use or to be incorporated into products for resale without written permission form Lifespan Technologies. Materials in this publication, as well as applications, methods and use, may be covered by one or more U.S. or foreign patents or patents pending. We welcome inquiries about licensing the use of our trademarks and technologies at info@lifespantech.com.