



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Lifespan Technologies
PO Box 3352
Salt Lake City, UT 84110
Tel: 801-971-1818 Fax: 801-464-6116
info@lifespantech.com

Product No: **S-7500**
Liquid Stable Conjugate
Approximate Dynamic Range
Dextran 10 kDa 3 -1000 µg/mL
Dextran 40 - 150 kDa 0.3 -100 µg/mL

Dextran ELISA Kit

INTENDED USE: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT INTENDED FOR CLINICAL OR DIAGNOSTIC USE.

Kit includes:

Coated 96-well plate
Detector -Enzyme Conjugate Solution
TMB Solution
Stop Solution
Wash Concentrate 10X, (dilute 1 part plus 9 parts water)


Researcher must provide:

Pipettes (8 Channel Multipipettor is recommended)
Absorbance microplate reader
Dextran Standards
Reagent Grade Water
Plate Sealer

Storage and Stability

Kit can be stored unopened at 4°C for up to six months. The Detector-Enzyme Conjugate Solution and the TMB solution should be protected from light.

Background



Dextran is a complex, branched glucan polysaccharide made of many glucose molecules composed of chains of varying lengths. The straight chain consists of α -1,6 glycosidic linkages between glucose molecules, while branches begin from α -1,4 linkages (and in some cases, α -1,2 and α -1,3 linkages as well). Dextran is synthesized from sucrose by certain lactic-acid bacteria, the best known being *Leuconostoc mesenteroides* and *Streptococcus mutans*. It is used medicinally as an antithrombotic (anti-platelet), to reduce blood viscosity and as a volume expander in anemia. It is used as a masking agent for some doping agents utilized in athletic performance enhancement. Dextran is damaging to the sugar industry for several reasons and should be minimized. It inhibits sucrose crystal formation which reduces product yield. It coats the surfaces it touches with a hard precipitate that can harm machinery for sugar producers and their customers. Dextran can cause product perception problems in food, beverage and liqueur.

The Dextran ELISA product number S-7500 is a quantitative enzyme-linked assay designed for the *in vitro* measurement of Dextran levels in fluids such as buffer or diluted juice (or syrup or molasses) from sugar cane or sugar beets. This assay measures Dextran directly using a Dextran antibody which has been conjugated to HRP.

The Dextran ELISA is a competitive assay in which the colorimetric signal is inversely proportional to the amount of Dextran present in the sample. Samples to be assayed are first mixed with the detector-enzyme conjugate in the wells of the coated plate. Dextran in the sample competes with Dextran bound to the plate for binding of the detector-enzyme conjugate. The bound peroxidase is reacted with tetramethylbenzidine (TMB) to quantitate the assay. The concentration of Dextran in the sample is determined using a standard curve of known amounts of Dextran. Parts per million and micrograms per mL are interchangeable. (1 PPM = 1 µg/mL)


Reagent Preparation

1X Wash Buffer: Make a 1:10 dilution of 10X Wash Buffer in distilled or deionized water

Sample Dilutions:

Samples can be assayed neat for most sensitive testing. If dilution is required, dilute with TBS.

Dextran Standards: Make dilutions of the appropriate MW Dextran standard using reagent grade water or TBS to obtain standards of 0.3, 1, 3, 10, 30 and 100µg/mL. **Standardization should be performed using Dextran that is the same Dextran type contained in your unknowns.** If the molecular weight of your samples is unknown, be advised that very small dextrans react less with our antibody, resulting in lower values.



Dextran ELISA

Assay Procedure

1. Set up the Dextran ELISA plate as illustrated above. We suggest the Dextran standard dilution series be run in triplicate for best results. Add **50µl** Standards and samples into corresponding wells. Add **50µl** of Detector-Enzyme conjugate to all wells except the Blank wells. Mix well. Cover plate and incubate for 30 minutes at room temperature. A rotator is highly recommended if available, as constant mixing significantly improves precision.
2. Discard the solution and wash the wells four times with 300 µL per well of 1X Wash Buffer. An automated plate washer is recommended if available. After washing, immediately proceed to the next step. Do not delay in removing wash buffer from the wells. Do not allow the plate to dry.
3. Add **100 µL** of TMB to all wells. Incubate the plate in the dark at room temperature for 10 - 60 minutes waiting for the zero Dextran wells to develop to a medium to dark blue color. Watch for color development and DO NOT overdevelop.
4. Add **50 µL** Stop Solution which will change the color from blue to yellow.
5. Immediately measure the absorbance of each well at 450 nm in an absorbance microplate reader.
6. Calculate the binding percentage for each sample using the formula:

$$[A_{450}(\text{Sample}) - A_{450}(\text{Blank})] / [A_{450}(\text{Zero Dextran}) - A_{450}(\text{Blank})] \times 100 = \% \text{ Binding}$$

Using linear or nonlinear regression, plot a standard curve of percent binding versus concentration of Dextran standards. We usually use a quadratic fit. Determine Dextran levels of unknowns by comparing their percentage binding relative to the standard curve. Dextran can be estimated by comparing the values from the wells containing unknowns to the values in the standard curve.

7. If samples were diluted, adjust results multiplying by the dilution factor.

Lifespan Technologies products are sold for research and development purposes only and are not for diagnostic use or to be incorporated into products for resale without written permission from Lifespan Technologies. Materials in this publication, as well as applications, methods and use, may be covered by one or more U.S. or foreign patents or patents pending. We welcome inquiries about licensing the use of our trademarks and technologies at info@lifespantech.com.