



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Ro 67-7476

Cat. No.:	HY-100403		
CAS No.:	298690-60-5		
Molecular Formula:	C ₁₇ H ₁₈ FNO ₂ S		
Molecular Weight:	319.39		
Target:	mGluR		
Pathway:	GPCR/G Protein; Neuronal Signaling		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 40 mg/mL (125.24 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		3.1310 mL	15.6548 mL	31.3097 mL
	5 mM		0.6262 mL	3.1310 mL	6.2619 mL
	10 mM		0.3131 mL	1.5655 mL	3.1310 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: ≥ 2.5 mg/mL (7.83 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
 Solubility: ≥ 2.5 mg/mL (7.83 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Ro 67-7476 is a potent positive allosteric modulator of mGluR₁ and potentiates glutamate-induced calcium release in HEK293 cells expressing rat mGluR1a with an EC₅₀ of 60.1 nM^{[1][2]}. Ro 67-7476 is a potent P-ERK1/2 agonist and activates ERK1/2 phosphorylation in the absence of exogenously added glutamate (EC₅₀=163.3 nM)^[3].

IC₅₀ & Target

mGluR1a
 60.1 nM (EC₅₀)

In Vitro

In the Purkinje cells of rat cerebellar slices, Ro 67-7476 increases the amplitude of mGluR1 excitatory postsynaptic potentials (EPSCs) evoked by 2,3-dihydroxy-6-nitro-7-sulfamoylbenzoquinoxaline, picrotoxin, or AP5^[3].

Ro 67-7476 activates ERK1/2 phosphorylation in the absence of exogenously added glutamate ($EC_{50}=163.3$ nM). The EC_{50} value of full P-ERK1/2 activation for Ro 67-7476 are nearly identical to the EC_{50} for calcium mobilization potentiation [3].

Ro 67-7476 increases basal cAMP production approximately by 8%. It potentiated threshold responses to glutamate in the cAMP accumulation assay, with an EC_{50} value of $17.7 \mu M$ [3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Int J Biol Sci. 2022 Jan 1;18(2):473-490.
- Int Immunopharmacol. 2022 Aug 20;111:109171.
- Mol Neurobiol. 2023 Sep 11.
- Pharmaceuticals. 2022 Aug 20;15(8):1027.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. F Knoflach, et al. Positive allosteric modulators of metabotropic glutamate 1 receptor: characterization, mechanism of action, and binding site. Proc Natl Acad Sci U S A. 2001 Nov 6;98(23):13402-7
- [2]. Kamondanai Hemstapat, et al. A novel class of positive allosteric modulators of metabotropic glutamate receptor subtype 1 interact with a site distinct from that of negative allosteric modulators. Mol Pharmacol. 2006 Aug;70(2):616-26.
- [3]. Douglas J Sheffler, et al. Allosteric potentiators of metabotropic glutamate receptor subtype 1a differentially modulate independent signaling pathways in baby hamster kidney cells. Neuropharmacology. 2008 Sep;55(4):419-27

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA