



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Splitomicin

Cat. No.:	HY-100585
CAS No.:	5690-03-9
Molecular Formula:	C ₁₃ H ₁₀ O ₂
Molecular Weight:	198.22
Target:	HDAC
Pathway:	Cell Cycle/DNA Damage; Epigenetics
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (504.49 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		5.0449 mL	25.2245 mL	50.4490 mL
		5 mM		1.0090 mL	5.0449 mL	10.0898 mL
		10 mM		0.5045 mL	2.5224 mL	5.0449 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (10.49 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (10.49 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (10.49 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Splitomicin (Splitomycin) is a selective Sir2p inhibitor. Splitomicin inhibits NAD ⁺ -dependent HDAC activity of Sir2 protein. Splitomicin induces dose-dependent inhibition of HDAC in the yeast extract with an IC ₅₀ of 60 μM ^[1] .
IC ₅₀ & Target	Sir2p 60 μM (IC ₅₀)
In Vitro	Splitomicin (10-333 μM; 24 hours) elicits antiproliferative effects in MCF-7 and H1299 cells in a dose-dependent manner in

colony formation assay. Splitomicin (33 μ M) fails to decrease the number of colonies, but Splitomicin (100 and 333 μ M) effectively inhibits colony formation in MCF-7 and H1299 cells^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Proliferation Assay^[2]

Cell Line:	Human breast cancer MCF-7 and lung cancer H1299 cells
Concentration:	10, 33, 100, and 333 μ M
Incubation Time:	24 hours
Result:	Inhibited colony formation in a dose-dependent manner.

In Vivo

Splitomicin (80 mg/kg with an intraperitoneal injection every 24 h for 5 days, in mice) enhances tissue factor (TF) activity in the arterial vessel wall and accelerates carotid artery thrombus formation in a photochemical injury model^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	C57BL/6 mice aged 12-14 weeks weighing on average 27 g ^[3]
Dosage:	80 mg/kg
Administration:	Intraperitoneal injection every 24 h for 5 days
Result:	Increased TF activity in mouse carotid artery as compared with the controls.

REFERENCES

[1]. Bedalov A, et al. Identification of a small molecule inhibitor of Sir2p. Proc Natl Acad Sci U S A. 2001 Dec 18;98(26):15113-8.

[2]. Breitenstein A, et al. Sirt1 inhibition promotes in vivo arterial thrombosis and tissue factor expression in stimulated cells. Cardiovasc Res. 2011 Feb 1;89(2):464-72.

[3]. Ota H, et al. Sirt1 inhibitor, Sirtinol, induces senescence-like growth arrest with attenuated Ras-MAPK signaling in human cancer cells. Oncogene. 2006 Jan 12;25(2):176-85.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA