



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



AS1842856

Cat. No.:	HY-100596		
CAS No.:	836620-48-5		
Molecular Formula:	C ₁₈ H ₂₂ FN ₃ O ₃		
Molecular Weight:	347.38		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 5 mg/mL (14.39 mM); ultrasonic and warming and heat to 60°C)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.8787 mL	14.3935 mL	28.7869 mL
	5 mM		0.5757 mL	2.8787 mL	5.7574 mL
	10 mM		0.2879 mL	1.4393 mL	2.8787 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 6% HP-β-CD in saline
Solubility: 10 mg/mL (28.79 mM); Suspension solution; Need ultrasonic
2. Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 1.67 mg/mL (4.81 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: 1 mg/mL (2.88 mM); Suspended solution; Need ultrasonic
4. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 1 mg/mL (2.88 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

AS1842856, a specific Foxo1 inhibitor (IC₅₀=30 nM), potently suppresses autophagy^[1]. AS1842856 only reduces the activity of FoxO1 by binding with it, without affecting its transcription and protein expression^[2].

IC₅₀ & Target

IC₅₀: 30 nM (Foxo1)^[1]

In Vitro

AS1842856 potently inhibits human Foxo1 transactivation and reduces glucose production through the inhibition of glucose-6 phosphatase and phosphoenolpyruvate carboxykinase mRNA levels in a rat hepatic cell line^[1]. After AS1842856

treatment, there is no significant difference in the protein expression of p-FoxO1 and FoxO1 compared with the control group, but the expression of p-Akt is decreased compared with the control group^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Oral administration of AS1842856 to diabetic db/db mice leads to a drastic decrease in fasting plasma glucose level via the inhibition of hepatic gluconeogenic genes, whereas administration to normal mice has no effect on the fasting plasma glucose level. Treatment with AS1842856 also suppresses an increase in plasma glucose level caused by pyruvate injection in both normal and db/db mice^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

Rat hepatoma Fao cells are cultured in DMEM with 5.5 mM glucose and 10% FBS. Glucose production rate is measured using glucose CII-test reagent. In brief, after 18 h of treatment with AS1842856 at the indicated concentrations, the cells are washed three times with PBS. The cells are then incubated for 3 h at 37°C in 5% CO₂ in a glucose production buffer (glucose-free DMEM, pH 7.4, containing 20 mM sodium pyruvate, without phenol red)^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

AS1842856 is dissolved in 6% cyclodextrin for oral administration. Pyruvate or glucose tolerance tests are performed in male mice aged 7 to 9 weeks. Mice are orally administered either AS1842856 dissolved in 6% cyclodextrin or vehicle (6% cyclodextrin only) at three time points (8 AM, 6 PM, and 8 AM on the second day). Food is removed after initial dosing and withheld throughout the study (26-h fasting)^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell. 2024 Feb 1;187(3):624-641.e23.
- J Exp Med. 2023 Oct 2;220(10):e20221653.
- Exp Mol Med. 2021 Sep;53(9):1307-1318.
- Theranostics. 2021 Jul 25;11(18):8624-8639.
- Redox Biol. 2022 Jun;52:102311.

See more customer validations on www.MedChemExpress.com

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA