



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Pan-RAS-IN-1

Cat. No.:	HY-101295
CAS No.:	1835283-94-7
Molecular Formula:	C <sub>36</sub> H <sub>41</sub> Cl <sub>2</sub> F <sub>3</sub> N <sub>6</sub> O <sub>2</sub>
Molecular Weight:	717.65
Storage:	<div> Powder -20°C 3 years </div> <div> 4°C 2 years </div> <div> In solvent -80°C 2 years </div> <div> -20°C 1 year </div>



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 50 mg/mL (69.67 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.3934 mL	6.9672 mL	13.9344 mL
	5 mM		0.2787 mL	1.3934 mL	2.7869 mL
	10 mM		0.1393 mL	0.6967 mL	1.3934 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: ≥ 2.5 mg/mL (3.48 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
 Solubility: ≥ 2.5 mg/mL (3.48 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
 Solubility: ≥ 2.5 mg/mL (3.48 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Pan-RAS-IN-1 is a pan-Ras inhibitor that disrupts the interaction of Ras proteins and their effectors.

#### In Vitro

Pan-RAS-IN-1 binds to KRas<sup>G12D</sup>-GppNHp with an affinity less than 20 μM. Pan-RAS-IN-1 binds to Ras proteins and exhibits lethality in cells partially dependent on expression of Ras proteins. The potency of pan-RAS-IN-1 correlates with the degree of dependency on the mutated isoform over a 5-fold concentration range. At some concentrations, pan-RAS-IN-1 is cytostatic, possibly due to pan-RAS inhibition. Pan-RAS-IN-1 is evaluated in primary T cell acute lymphoblastic leukemia (T-ALL) cells. Selective lethality is observed, with mutant NRAS cells retaining only 20%-40% viability after 5 μM treatment<sup>[1]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## In Vivo

Pan-RAS-IN-1 administration results in inhibition of tumor growth over 15 days of treatment. Pan-RAS-IN-1-treated mice exhibits decreased tumor pERK levels compared with vehicle treated mice. A modest increase in cleaved caspase-3 is also observed, showing that in this model, pan-RAS-IN-1 has the capacity to induce caspase activation<sup>[1]</sup>.  
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

### Cell Assay <sup>[1]</sup>

For 384-well cancer cell viability assays, cells are trypsinized, counted, and seeded into 384-well plates at 1,000 cells/well. After 16 hr, pan-RAS-IN-1 (from 10 mM stocks in DMSO) are arrayed in an 8-point or 16-point dilution series in 384-well polypropylene plates. Compound solutions are transferred at a 1:5 dilution into the assay plates. After 48 hr, a 50% Alamar blue solution is added to a final concentration of 10% Alamar blue. After 6 hr of incubation, fluorescence intensity is determined at 535 and 590 nm<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### Animal Administration <sup>[1]</sup>

Mice: Mice tumor Xenograft are dosed with 180 mg/kg pan-RAS-IN-1 orally (12 mg/mL, 10% DMSO, pH 4), vehicle orally, or by a combination of i.p. and i.v. injections at 30 mg/kg (4 mg/mL, 5% DMSO in HBSS at pH 4). Over 14 d, mice receive a total of 10 doses of pan-RAS-IN-1 or vehicle orally, or six i.p. injections and 4 i.v. injections. Tumor size is measured by electronic caliper every 2 d and calculated<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- Cell Rep. 2021 Jul 6;36(1):109315.
- Cell Chem Biol. 2021 May 4;S2451-9456(21)00206-3.
- Antiviral Res. 2021 May 4;105082.
- Dis Model Mech. 2022 Feb 1;15(2):dmm049093.
- Neurooncol Adv. 31 December 2021.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA