



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NPS-2143 hydrochloride

Cat. No.:	HY-10171
CAS No.:	324523-20-8
Molecular Formula:	C ₂₄ H ₂₆ Cl ₂ N ₂ O ₂
Molecular Weight:	445.38
Target:	CaSR
Pathway:	GPCR/G Protein
Storage:	4°C, sealed storage, away from moisture
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 100 mg/mL (224.53 mM)

H₂O : 1.85 mg/mL (4.15 mM; ultrasonic and warming and heat to 60°C)

* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
	Concentration				
	1 mM	2.2453 mL	11.2264 mL	22.4527 mL	
	5 mM	0.4491 mL	2.2453 mL	4.4905 mL	
	10 mM	0.2245 mL	1.1226 mL	2.2453 mL	

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline

Solubility: ≥ 2.5 mg/mL (5.61 mM); Clear solution

2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)

Solubility: ≥ 2.5 mg/mL (5.61 mM); Clear solution

3. Add each solvent one by one: 10% DMSO >> 90% corn oil

Solubility: ≥ 2.5 mg/mL (5.61 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	NPS-2143 hydrochloride (SB-262470A hydrochloride), an orally active calcilytic agent, is a selective and potent calcium ion-sensing receptor (CaSR) antagonist. NPS-2143 hydrochloride (SB-262470A hydrochloride) blocks increases in cytoplasmic Ca ²⁺ concentrations (IC ₅₀ =43 nM) elicited by activating the Ca ²⁺ receptor in HEK 293 cells expressing the human Ca ²⁺ receptor [1][2].
In Vitro	NPS-2143 hydrochloride (SB-262470A hydrochloride) stimulates parathyroid hormone (PTH) secretion from bovine parathyroid cells with EC ₅₀ of 41 nM. Moreover, NPS-2143 hydrochloride also blocks the inhibitory effects of calcimimetic

	NPS R-467 on PTH secretion from bovine parathyroid cells and the inhibitory effects of extracellular Ca²⁺ on isoproterenol-stimulated increases in cyclic AMP formation^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	When infused intravenously in normal rats, NPS-2143 hydrochloride (SB-262470A hydrochloride) causes a rapid and large increase in plasma levels of PTH. Ca²⁺ receptor antagonists are termed calcilytics and NPS-2143 is the first substance (either atomic or molecular) shown to possess such activity^[1]. When administered together with an antiresorptive agent (estradiol), NPS 2143 causes an increase in trabecular bone volume and bone mineral density in osteopenic rats^[2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Phytomedicine. 2021, 153507.
- Acta Physiol. 2023 Jan 6;e13926.
- Food Funct. 2022 May 19.
- Int J Mol Sci. 2023 Mar 3.
- Front Pharmacol. 2022 Feb 23;13:816133.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Huang Y, Breitwieser GE. Rescue of calcium-sensing receptor mutants by allosteric modulators reveals a conformational checkpoint in receptor biogenesis. J Biol Chem. 2007 Mar 30;282(13):9517-25
- [2]. Marquis, Robert W.; Lago, Amparo M.; Callahan, James F.;Antagonists of the Calcium Receptor. 2. Amino Alcohol-Based Parathyroid Hormone Secretagogues. Journal of Medicinal Chemistry (2009), 52(21), 6599-6605.
- [3]. Yamamura A, Guo Q, Yamamura H, Zimnicka AM, Pohl NM, Smith KA, Fernandez RA, Zeifman A, Makino A, Dong H, Yuan JX.Enhanced Ca²⁺-sensing receptor function in idiopathic pulmonary arterial hypertension.Circ Res. 2012 Aug 3;111(4):469-81. Epub 2012 Jun 22.
- [4]. Nakajima S, Hira T, Hara H.Calcium-sensing receptor mediates dietary peptide-induced CCK secretion in enteroendocrine STC-1 cells.Mol Nutr Food Res. 2012 May;56(5):753-60.
- [5]. Davey AE, Leach K, Valant C, Conigrave AD, Sexton PM, Christopoulos A.Positive and negative allosteric modulators promote biased signaling at the calcium-sensing receptor.Endocrinology. 2012 Mar;153(3):1232-41. Epub 2011 Dec 30.
- [6]. Nemeth EF, et al. Calcilytic compounds: potent and selective Ca²⁺ receptor antagonists that stimulate secretion of parathyroid hormone. J Pharmacol Exp Ther. 2001 Oct;299(1):323-31.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA