



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Quin-2AM

Cat. No.:	HY-101902
CAS No.:	83104-85-2
Molecular Formula:	C ₃₈ H ₄₃ N ₃ O ₁₈
Molecular Weight:	829.76
Target:	Fluorescent Dye
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Quin-2AM is a fluorescent Ca ²⁺ chelator, with high affinity for calcium. Quin-2AM can specifically identify intracellular calcium ions, with high sensitivity, low cytotoxicity, increased AM acetylmethyl ester can enter the cell well, after being sheared by the intracellular esterase stay in the cell to bind to calcium ions, produce strong fluorescence ^[1] .
In Vitro	<ol style="list-style-type: none"> Preparation of Quin-2AM working solution <ol style="list-style-type: none"> Preparation of the stock solution Dissolve 1 mg Quin-2AM in 135 μL DMSO to obtain 10 mM of stock solution. Note: It is recommended to store the stock solution at -20℃ or -80℃ away from light and avoid repetitive freeze-thaw cycles. Preparation of Quin-2AM working solution Dilute the stock solution in HBSS to obtain 1-10 μM of working solution. Note: Please adjust the concentration of Quin-2AM working solution according to the actual situation. Cell staining <ol style="list-style-type: none"> Suspension cells (6-well plate) <ol style="list-style-type: none"> Centrifuge at 1000 g at 4℃ for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. The cell density is 1×10⁶/mL. Add 1 mL of working solution, and then incubate at room temperature for 5-30 minutes. Centrifuge at 400 g at 4℃ for 3-4 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. Resuspend cells with serum-free cell culture medium or PBS. Observation by fluorescence microscopy or flow cytometry. Adherent cells <ol style="list-style-type: none"> Culture adherent cells on sterile coverslips. Remove the coverslip from the medium and aspirate excess medium. Add 100 μL of working solution, gently shake it to completely cover the cells, and then incubate at room temperature for 5-30 minutes. Wash twice with medium, 5 minutes each time. Observation by fluorescence microscopy. <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>

REFERENCES

- [1]. Chu AJ, et al. Possible role of Marcks in the cellular modulation of monocytic tissue factor-initiated hypercoagulation. Br J Haematol. 2002 Aug;118(2):569-76.
- [2]. Chu AJ, et al. Possible role of Marcks in the cellular modulation of monocytic tissue factor-initiated hypercoagulation. Br J Haematol. 2002 Aug;118(2):569-76.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA