



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BETP

Cat. No.:	HY-103546
CAS No.:	1371569-69-5
Molecular Formula:	C ₂₀ H ₁₇ F ₃ N ₂ O ₂ S
Molecular Weight:	406.42
Target:	GCGR
Pathway:	GPCR/G Protein
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : 25 mg/mL (61.51 mM; Need ultrasonic)
H₂O : < 0.1 mg/mL (insoluble)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.4605 mL	12.3025 mL	24.6051 mL
	5 mM		0.4921 mL	2.4605 mL	4.9210 mL
	10 mM		0.2461 mL	1.2303 mL	2.4605 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

BETP is an agonist of glucagon-like peptide-1 (GLP-1) receptor, with EC₅₀s of 0.66 and 0.755 μM for human and rat GLP-1 receptor, respectively.

IC₅₀ & Target

EC₅₀: 0.66 μM (Human GLP-1 receptor), 0.755 μM (Rat GLP-1 receptor)^[1]

In Vitro

BETP is a GLP-1 receptor agonist, with EC₅₀s of 0.66 and 0.755 μM for human and rat GLP-1 receptor, respectively. BETP (Compound B) is inactive in cells expressing the GLP-2, GIP, PTH, or glucagon receptors. BETP (1-10 μM) enhances insulin secretion in normal and diabetic human islets. In addition, BETP in combination with GLP-1 shows additive effects on increasing GLP-1 receptor signaling^[1]. BETP increases the potency of oxyntomodulin by 10-fold (EC₅₀ of 80 pM). GLP-1 does

not change the potencies and efficacies of both oxyntomodulin and glucagon at the glucagon receptor. BETP (0-30 μ M) increases the binding affinity of oxyntomodulin for the GLP-1 receptor^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

BETP has insulinotropic effect in SD rats. BETP (10 mg/kg, jugular vein cannula) exhibits insulin secretagogue activity in the intravenous glucose tolerance test (IVGTT) model. BETP (10 mg/kg, i.v.)-treated rats need 20% higher glucose infusion rates and demonstrates higher plasma insulin levels in SD rat hyperglycemic clamp model^[1]. BETP (5 mg/kg) enhances oxyntomodulin-stimulated insulin secretion^[2].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Animal Administration ^[1]

Rats^[1]

The IVGTT studies are performed. Male SD rats are group-housed three per cage in polycarbonate cages with filter tops. Rats are maintained on a 12:12 h light-dark cycle (lights on at 6:00 a.m.) at 21°C and receive diet and deionized water ad libitum. Rats are fasted overnight and anesthetized with 60 mg/kg pentobarbital for the duration of the experiment. For glucose and compounds (BETP, etc.) administration, a catheter with a diameter of 0.84 mm is inserted into the jugular vein. For rapid blood collection, a larger catheter with 1.02-mm diameter is inserted into the carotid artery. Blood is collected for glucose and insulin levels at time 0, 2, 4, 6, 10, and 20 min after intravenous administration of the BETP which is immediately followed by an intravenous glucose bolus of 0.5 g/kg. Plasma levels of glucose and insulin are determined^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Sloop KW, et al. Novel small molecule glucagon-like peptide-1 receptor agonist stimulates insulin secretion in rodents and from human islets. Diabetes. 2010 Dec;59(12):3099-107.

[2]. Willard FS, et al. Small molecule allosteric modulation of the glucagon-like Peptide-1 receptor enhances the insulinotropic effect of oxyntomodulin. Mol Pharmacol. 2012 Dec;82(6):1066-73.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA