



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AMN082 free base

Cat. No.:	HY-103565A
CAS No.:	83027-13-8
Molecular Formula:	C ₂₈ H ₂₈ N ₂
Molecular Weight:	392.54
Target:	mGluR
Pathway:	GPCR/G Protein; Neuronal Signaling
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 12.5 mg/mL (31.84 mM; ultrasonic and warming and heat to 60°C)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.5475 mL	12.7376 mL	25.4751 mL
		5 mM		0.5095 mL	2.5475 mL	5.0950 mL
		10 mM		0.2548 mL	1.2738 mL	2.5475 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 1.25 mg/mL (3.18 mM); Suspended solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 1.25 mg/mL (3.18 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	AMN082 free base, a selective, orally active, and brain penetrant mGluR7 agonist, directly activates receptor signaling via an allosteric site in the transmembrane domain. AMN082 free base potently inhibits cAMP accumulation and stimulates GTPγS binding (EC ₅₀ values, 64-290 nM) at transfected mammalian cells expressing mGluR7. AMN082 free base shows selectivity over other mGluR subtypes and selected ionotropic glutamate receptors. Antidepressant effects ^{[1][2]} .
IC ₅₀ & Target	mGluR7 ^[1]
In Vitro	Preincubation of the synaptosomes with AMN082 (1 μM) for 10 min before 4-aminopyridine treatment efficiently inhibits the 4-aminopyridine-evoked release of glutamate, without altering the basal release of glutamate ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

AMN082 (6 mg/kg; p.o.) induces stress hormone increases in an mGluR7-dependent fashion in mGluR7^{+/+} mice (C57BL/6 genetic background)^[1].

AMN082 (1.25-5.0 mg/kg, i.p.; 30 min before every Cocaine injection during repeated drug administration or before Cocaine challenge) dose-dependently attenuates the development, as well as the expression of Cocaine locomotor sensitization^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Mitsukawa K, et al. A selective metabotropic glutamate receptor 7 agonist: activation of receptor signaling via an allosteric site modulates stress parameters in vivo. *Proc Natl Acad Sci U S A*. 2005;102(51):18712-18717.

[2]. Wang CC, et al. Metabotropic glutamate 7 receptor agonist AMN082 inhibits glutamate release in rat cerebral cortex nerve terminal. *Eur J Pharmacol*. 2018;823:11-18.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA