



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

L-Ascorbic acid 2-phosphate magnesium hydrate

Cat. No.:	HY-103701B
CAS No.:	1713265-25-8
Molecular Formula:	C ₆ H ₉ O ₉ P.xH ₂ O.3/2Mg
Target:	Phosphatase; Reactive Oxygen Species; Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease; Immunology/Inflammation; NF-κB
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro	H ₂ O : 5 mg/mL (ultrasonic and warming and heat to 60°C)
-----------------	--

BIOLOGICAL ACTIVITY

Description	L-ascorbic acid 2-phosphate (2-Phospho-L-ascorbic acid) magnesium hydrate is a long-acting vitamin C derivative that can stimulate collagen formation and expression. L-ascorbic acid 2-phosphate magnesium hydrate can be used as a culture medium supplement for the osteogenic differentiation of human adipose stem cells (hASCs). L-ascorbic acid 2-phosphate magnesium hydrate increases alkaline phosphatase (ALP) activity and expression of runx2A in hASCs during the osteogenic differentiation ^{[1][2][3]} .
--------------------	---

In Vitro	<p>L-Ascorbic acid 2-phosphate (0.1-1.5 mM; 2 to 3 weeks with medium exchange every 2 to 3 days) magnesium hydrate significantly stimulates cell growth, whereas addition of l-Ascorbic acid (Asc) achieves only weak growth stimulation. A combination of Asc-2P and bFGF significantly increases cell growth, but supplementation with EGF and/or insulin does not have any additional effect^[1].</p> <p>L-Ascorbic acid 2-phosphate (50 μM-250 μM) magnesium hydrate is needed for the effective osteogenic differentiation of human adipose stem cells (hASCs), and higher concentrations of AsA2-P results in increased runx2 expression and ALP activity. The highest proliferation, ALP activity and runx2 expression is achieved with 150 μM AsA2-P and 10 nM dexamethasone (Dex), and 250 μM AsA2-P and 5 nM Dex^[3].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
-----------------	--

Cell Proliferation Assay^[1]

Cell Line:	Human corneal endothelial cells (HCECs)
Concentration:	0.1 mM; 0.3 mM; 1.5 mM
Incubation Time:	2 to 3 weeks with medium exchange every 2 to 3 days
Result:	Stimulated HCEC cells growth.

REFERENCES

-
- [1]. Shima N, et al. Increased proliferation and replicative lifespan of isolated human corneal endothelial cells with L-ascorbic acid 2-phosphate. Invest Ophthalmol Vis Sci. 2011 Nov 7;52(12):8711-7.
- [2]. Kurata S, et al. Epidermal growth factor inhibits transcription of type I collagen genes and production of type I collagen in cultured human skin fibroblasts in the presence and absence of L-ascorbic acid 2-phosphate, a long-acting vitamin C derivative. J Biol Chem. 1991 May 25;266(15):9997-10003.
- [3]. Kyllönen L, et al. Effects of different serum conditions on osteogenic differentiation of human adipose stem cells in vitro. Stem Cell Res Ther. 2013 Feb 15;4(1):17.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA