



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



AH 6809

Cat. No.:	HY-10418
CAS No.:	33458-93-4
Molecular Formula:	C ₁₇ H ₁₄ O ₅
Molecular Weight:	298.29
Target:	Prostaglandin Receptor
Pathway:	GPCR/G Protein
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (83.81 mM; Need ultrasonic) H ₂ O : < 0.1 mg/mL (insoluble)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	3.3524 mL	16.7622 mL	33.5244 mL
		5 mM	0.6705 mL	3.3524 mL	6.7049 mL
		10 mM	0.3352 mL	1.6762 mL	3.3524 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (8.38 mM); Suspended solution; Need ultrasonic				
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: 2.5 mg/mL (8.38 mM); Suspended solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	AH 6809 is an antagonist of EP and DP receptor, with K _i s of 1217, 1150, 1597, and 1415 nM for the cloned human EP ₁ , EP ₂ , EP ₃ -III, and DP receptor respectively. AH 6809 has a K _i of 350 nM for mouse EP ₂ receptor ^{[1][2][3]} .
In Vitro	AH 6809 (1 μM; 30 min) inhibits T. serrulatus venom (TsV)-induced and PGE ₂ -amplified IL-1β and cAMP production in macrophages ^[4] . AH 6809 (30-300 μM) antagonizes the anti-aggregatory activity of PGD ₂ in whole blood, with an apparent pA ₂ of 5.35 ^[5] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	AH 6809 (5 mg/kg; i.p.) decreases TsV-induced mortality, PGE ₂ and IL-1β production and neutrophil infiltration in the lungs

of mice^[4].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Immunol. 2020 Jan;347:104025.
- Neurol Int. 2022, 14(1), 11-33.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Abramovitz M, et, al. The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. Biochim Biophys Acta. 2000 Jan 17;1483(2):285-93.
- [2]. Kiriya M, et, al. Ligand binding specificities of the eight types and subtypes of the mouse prostanoid receptors expressed in Chinese hamster ovary cells. Br J Pharmacol. 1997 Sep;122(2):217-24.
- [3]. Woodward DF, et, al. 6-Isopropoxy-9-oxoxanthene-2-carboxylic acid (AH 6809), a human EP2 receptor antagonist. Biochem Pharmacol. 1995 Nov 9;50(10):1731-3.
- [4]. Zoccal KF, et, al. Opposing roles of LTB4 and PGE2 in regulating the inflammasome-dependent scorpion venom-induced mortality. Nat Commun. 2016 Feb 23;7:10760.
- [5]. Keery RJ, et, al. AH6809, a prostaglandin DP-receptor blocking drug on human platelets. Br J Pharmacol. 1988 Jul;94(3):745-54.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA