



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Darapladib

Cat. No.:	HY-10521
CAS No.:	356057-34-6
Molecular Formula:	C ₃₆ H ₃₈ F ₄ N ₄ O ₂ S
Molecular Weight:	666.77
Target:	Phospholipase; Apoptosis
Pathway:	Metabolic Enzyme/Protease; Apoptosis
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 100 mg/mL (149.98 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.4998 mL	7.4988 mL	14.9977 mL
	5 mM		0.3000 mL	1.4998 mL	2.9995 mL
	10 mM		0.1500 mL	0.7499 mL	1.4998 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (3.75 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (3.75 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (3.75 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Darapladib (SB-480848) is an orally active, selective and reversible Lp-PLA2 inhibitor (IC₅₀=0.25 nM). Darapladib can trigger irreversible actions on glioma cell apoptosis and induce cycle arrest. Darapladib can be used in the study of atherosclerosis and cancer^{[1][2][3][4]}.

IC₅₀ & Target

IC₅₀: 0.25 nM (Lp-PLA₂)^[1]

In Vitro

Darapladib (5 μ M; 6, 12 h) induces cell cycle arrest in glioma cells (C6 glioma cells and U251MG cells)^[2].

Darapladib (5 μ M; 3, 6 h) triggers cell apoptosis in glioma cells^[2].

Darapladib (5 μ M; 5, 15, 30, 60 and 90 min) induces an increase in phosphorylation of ERK1/2 proteins in glioma cells^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Apoptosis Analysis^[2]

Cell Line:	C6 glioma cells and U251MG cells.
Concentration:	5 μ M
Incubation Time:	3, 6 h
Result:	Triggered cell apoptosis in glioma cells.

Cell Cycle Analysis^[2]

Cell Line:	C6 glioma cells and U251MG cells.
Concentration:	5 μ M
Incubation Time:	6, 12 h
Result:	Induced cell cycle arrest in glioma cells.

Western Blot Analysis^[2]

Cell Line:	C6 glioma cells and U251MG cells.
Concentration:	5 μ M
Incubation Time:	5, 15, 30, 60 and 90 min
Result:	Induced an increase in phosphorylation of ERK1/2 proteins, but reduced AKT phosphorylation in glioma cells.

In Vivo

Darapladib (50 mg/kg; p.o.; once daily for 6 weeks) significantly inhibits serum Lp-PLA2 activity in LDLR-deficient mice^[3].

Darapladib (50 mg/kg; p.o.; once daily for 6 weeks) decreases serum hs-CRP and IL-6 levels^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male homozygous LDLR-deficient mice (C57/BL6 genetic background) ^[3] .
Dosage:	50 mg/kg
Administration:	Oral administration; once daily for 6 weeks.
Result:	Significantly inhibited activity of serum Lp-PLA2.

CUSTOMER VALIDATION

- J Med Chem. 2016 May 26;59(10):5115-20.
- Eur J Pharmacol. 2020 Nov 15;887:173559.
- Sci Rep. 2017 Oct 3;7(1):12628.
- RSC Adv. 2017, 7(83):52762-52771.

-
- Università Vita-Salute San Raffaele. 2022 Apr 08. 34.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Wang YJ, et al. The selective lipoprotein-associated phospholipase A2 inhibitor darapladib triggers irreversible actions on glioma cell apoptosis and mitochondrial dysfunction. *Toxicol Appl Pharmacol*. 2020 Sep 1;402:115133.
- [2]. Riley RF, et al. Darapladib, a reversible lipoprotein-associated phospholipase A2 inhibitor, for the oral treatment of atherosclerosis and coronary artery disease. *IDrugs*. 2009 Oct;12(10):648-55.
- [3]. Blackie JA, et al. The identification of clinical candidate SB-480848: a potent inhibitor of lipoprotein-associated phospholipase A2. *Bioorg Med Chem Lett*. 2003 Mar 24;13(6):1067-70.
- [4]. Hu MM, et al. The inhibition of lipoprotein-associated phospholipase A2 exerts beneficial effects against atherosclerosis in LDLR-deficient mice. *Acta Pharmacol Sin*. 2011 Oct;32(10):1253-1258.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA