



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Nelotanserine

Cat. No.:	HY-10559
CAS No.:	839713-36-9
Molecular Formula:	C <sub>18</sub> H <sub>15</sub> BrF <sub>2</sub> N <sub>4</sub> O <sub>2</sub>
Molecular Weight:	437.24
Target:	5-HT Receptor
Pathway:	GPCR/G Protein; Neuronal Signaling
Storage:	<div> Powder -20°C 3 years </div> <div> 4°C 2 years </div> <div> In solvent -80°C 2 years </div> <div> -20°C 1 year </div>



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 32 mg/mL (73.19 mM)  
 \* "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		2.2871 mL	11.4354 mL	22.8707 mL
	5 mM		0.4574 mL	2.2871 mL	4.5741 mL
	10 mM		0.2287 mL	1.1435 mL	2.2871 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

Description	Nelotanserine is a potent 5-HT <sub>2A</sub> inverse agonist, a moderately potent 5-HT <sub>2C</sub> partial inverse agonist and a weak 5-HT <sub>2B</sub> inverse agonist, with IC <sub>50</sub> s of 1.7, 79, 791 nM in IP accumulation assays, respectively.		
IC <sub>50</sub> & Target	5-HT <sub>2A</sub> Receptor 1.7 nM (IC <sub>50</sub> )	5-HT <sub>2C</sub> Receptor 79 nM (IC <sub>50</sub> )	5-HT <sub>2B</sub> Receptor 791 nM (IC <sub>50</sub> )
In Vitro	<p>Results from IP accumulation assays suggest that Nelotanserine is a potent 5-HT<sub>2A</sub> full inverse agonist (IC<sub>50</sub>=1.7 nM), a moderately potent 5-HT<sub>2C</sub> partial inverse agonist (IC<sub>50</sub>=79 nM) (maximal response was 62% of the response obtained for the reference inverse agonist clozapine), and a weak 5-HT<sub>2B</sub> inverse agonist (IC<sub>50</sub>=791 nM). Nelotanserine displays high affinity for recombinant human 5-HT<sub>2A</sub> receptors (K<sub>i</sub>=0.35 nM), moderate affinity for human 5-HT<sub>2C</sub> receptors (K<sub>i</sub>=100 nM), and low affinity for human 5-HT<sub>2B</sub> receptors (2000 nM) stably expressed in HEK293 cells. The results suggest that Nelotanserine has a 262-fold higher affinity for human 5-HT<sub>2A</sub> than 5-HT<sub>2C</sub> receptors and a 6610-fold higher affinity for human 5-HT<sub>2A</sub> than 5-HT<sub>2B</sub> receptors<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>		

## In Vivo

Each compound is tested in a minimum of five rats by oral gavage with administration occurring in the middle of the inactive period, 6 h after light onset. The delta power during non-REM sleep (NREMS) is significantly different between all the analogues tested and the vehicle control. Nelotanserin (Compound 39) produces significant increases in delta power that persist for the first 4 h following dosing. Significant differences are found, however, in NREMS bout length. Nelotanserin significantly increases NREMS bout length during the first hour following dosing, and 3 does so during the second hour. In conjunction with this increased NREM bout duration, the number of NREM bouts decrease during the first hour for Nelotanserin ( $p < 0.01$ ) as well as for compound 15 ( $p < 0.05$ )<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## CUSTOMER VALIDATION

- ACS Chem Neurosci. 2019 Nov 20;10(11):4476-4491.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Al-Shamma HA et al. Nelotanserin, a novel selective human 5-hydroxytryptamine<sub>2A</sub> inverse agonist for the treatment of insomnia. J Pharmacol Exp Ther. 2010 Jan;332(1):281-90.

[2]. Teegarden BR et al. Discovery of 1-[3-(4-bromo-2-methyl-2H-pyrazol-3-yl)-4-methoxyphenyl]-3-(2,4-difluorophenyl)urea (nelotanserin) and related 5-hydroxytryptamine<sub>2A</sub> inverse agonists for the treatment of insomnia. J Med Chem. 2010 Mar 11;53(5):1923-36.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA