



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SY-640

Cat. No.:	HY-106947
CAS No.:	168705-70-2
Molecular Formula:	C ₁₁ H ₁₃ NO ₃
Molecular Weight:	207.23
Target:	Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	Pure form -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (482.56 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM	4.8256 mL	24.1278 mL	48.2556 mL	
		5 mM	0.9651 mL	4.8256 mL	9.6511 mL	
	10 mM	0.4826 mL	2.4128 mL	4.8256 mL		
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (12.06 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	SY-640 is an Acetamide derivative and has potent hepatoprotective effect. SY-640 reduces Propionibacterium acnes and Lipopolysaccharide-induced liver injury in mice ^{[1][2]} .
In Vitro	SY-640 inhibits the number of liver-infiltrating cells and attenuates the increased expression of leukocyte function-associated antigen-1 on these cells ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	SY-640 (Oral; 150 mg/kg; once daily for 7 days) significantly inhibits Propionibacterium acnes and Lipopolysaccharide-induced liver injury, but a single administration is without effect ^[1] . SY-640 (p.o.; 150 mg/kg; once daily for three days) significantly increases the liver microsomal cytochrome P-450 content and aminopyrine demethylase activity in mice. The hepatic microsomal aminopyrine demethylase activity is obviously inhibited two hours after oral administration of SY-640 ^[2] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Y Tanaka, et al. Hepatoprotective effect of SY-640, a novel acetamide derivative, on Propionibacterium acnes and lipopolysaccharide-induced liver injury in mice. Arch Int Pharmacodyn Ther. Mar-Apr 1995;329(2):319-30.
- [2]. P F Li, et al. [Inhibitory effect of 2-(N-acetyl-methyl amino)-3',4'-methylenedioxyacetyl-aminophene(SY-640) on covalent binding of carcinogenic benzo(a)pyrene with mouse hepatocyte nuclear DNA]. Yao Xue Xue Bao. 1997 Sep;32(9):663-8.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA