



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



SGN-2FF

Cat. No.:	HY-107366
CAS No.:	2089647-47-0
Molecular Formula:	C ₆ H ₁₁ FO ₄
Molecular Weight:	166.15
Target:	Others
Pathway:	Others
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : 50 mg/mL (300.93 mM; Need ultrasonic)
H₂O : 36.67 mg/mL (220.70 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		6.0187 mL	30.0933 mL	60.1866 mL
	5 mM		1.2037 mL	6.0187 mL	12.0373 mL
	10 mM		0.6019 mL	3.0093 mL	6.0187 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

1. Add each solvent one by one: PBS
Solubility: 50 mg/mL (300.93 mM); Clear solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Description	SGN-2FF is a potent and orally active inhibitor of fucosylation, directly inhibits fucosyltransferase activity. SGN-2FF possesses antitumor activity ^[1] .
IC ₅₀ & Target	Fucosyltransferase ^[1]
In Vitro	SGN-2FF (2-Fluorofucose) is an inhibitor of fucosylation, inhibits cellular fucosylation by depleting the fucosylation substrate GDP-fucose, and by direct inhibition of fucosyltransferases, and leads to the production of afucosylated glycoproteins including antibodies. SGN-2FF activates human T cells in an antigen-dependent manner ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	SGN-2FF exhibits antitumor activity in multiple mouse tumor models, showing substantial tumor growth delay. SGN-2FF

elevates the protective effect of a lymphoma vaccine in a syngeneic mouse model^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Stephen C. Alley, et al. Abstract DDT02-02: SGN-2FF: A novel small molecule inhibitor of fucosylation with preclinical antitumor activity through multiple immune mechanisms. Cancer Res 2017;77(13 Suppl).

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA