



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

σ1 Receptor antagonist-1

Cat. No.:	HY-10815
CAS No.:	1204401-49-9
Molecular Formula:	C ₂₁ H ₂₃ NOS
Molecular Weight:	337.48
Target:	Sigma Receptor; Apoptosis
Pathway:	Neuronal Signaling; Apoptosis
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (296.31 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.9631 mL	14.8157 mL	29.6314 mL
		5 mM		0.5926 mL	2.9631 mL	5.9263 mL
		10 mM		0.2963 mL	1.4816 mL	2.9631 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.41 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.41 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	σ1 Receptor antagonist-1 is a highly potent and selective sigma 1 receptor antagonist (pK _i =10.28). σ1 Receptor antagonist-1 inhibits cell growth, arrests cell cycle at G0/G1 phase and induces apoptosis of MCF-7/ADR cells ^[1] .
IC ₅₀ & Target	Sigma 1 Receptor
In Vitro	σ1 Receptor antagonist-1 (compound 9) (10 μM; 48 hours) inhibits cell growth, arrests cell cycle at G0/G1 phase and induces apoptosis of MCF-7/ADR cells ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. Cell Cycle Analysis ^[1]

Cell Line:	MCF-7/ADR, MCF-7 cells
Concentration:	10 μ M
Incubation Time:	48 hours
Result:	Increased the number of cells in G0/G1 (27.8%) and decreased those in S phase (46.7%), in a sigma 1-dependent manner. It selectively affects the high sigma 1 receptor expressing MCF-7/ADR but not the low sigma 1 receptor expressing MCF-7 cells.

In Vivo

Pretreatment with σ 1 Receptor antagonist-1 (1 mg/kg; s.c.) significantly increases the anti-nociceptive effect produced by morphine over the entire time course starting at 60 min^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male CD-1 mice ^[1]
Dosage:	1 mg/kg
Administration:	S.c.
Result:	Significantly increased the anti-nociceptive effect produced by morphine over the entire time course starting at 60 min.

CUSTOMER VALIDATION

- Cells. 2023 Jan 3;12(1):197.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Piergentili A, et al. Novel highly potent and selective sigma 1 receptor antagonists related to spipethiane. J Med Chem. 2010;53(3):1261-1269.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA