



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GNF-7

Cat. No.:	HY-10943
CAS No.:	839706-07-9
Molecular Formula:	C ₂₈ H ₂₄ F ₃ N ₇ O ₂
Molecular Weight:	547.53
Target:	Bcr-Abl; Ack1
Pathway:	Protein Tyrosine Kinase/RTK
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 33 mg/mL (60.27 mM)
 * "≥" means soluble, but saturation unknown.

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.8264 mL	9.1319 mL	18.2638 mL
	5 mM		0.3653 mL	1.8264 mL	3.6528 mL
	10 mM		0.1826 mL	0.9132 mL	1.8264 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
 Solubility: ≥ 2 mg/mL (3.65 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
 Solubility: ≥ 2 mg/mL (3.65 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

GNF-7 is a multikinase inhibitor. GNF-7 is a Bcr-Abl inhibitor, with IC₅₀s of 133 nM and 61 nM for Bcr-Abl^{WT} and Bcr-Abl^{T315I}, respectively. GNF-7 also possesses inhibitory activity against both ACK1 (activated CDC42 kinase 1) and GCK (germinal center kinase) with IC₅₀s of 25 nM and 8 nM, respectively. GNF-7 can be used for the research of hematologic malignancies^[1] [2][3].

IC₅₀ & Target

IC₅₀: 133 nM (Bcr-Abl^{WT})^[1], 61 nM (Bcr-Abl^{T315I})^[1], 25 nM (ACK1)^[3], 8 nM (GCK)^[3]

In Vitro

GNF-7 potently inhibits wild-type Bcr-Abl (IC₅₀<5 nM) and Bcr-Abl mutants such as T315I (IC₅₀=11 nM), G250E (IC₅₀<5 nM), E255V (IC₅₀=10 nM), F317L (IC₅₀<5 nM) and M351T (IC₅₀<5 nM) in cellular assays^[2].

GNF-7 (1 μ M; 2 hours) suppresses AKT/mTOR signaling and GCK downstream^[3].

GNF-7 (1 μ M; 24 hours) induces of apoptosis and cell cycle arrest in NRAS mutant cell lines^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Western Blot Analysis^[3]

Cell Line:	Ba/F3-NRAS-G12D cells, OCI-AML3 cells
Concentration:	1 μ M
Incubation Time:	2 hours
Result:	Caused a decreased level of phosphorylation of p70S6K1, AKT (S473), JNK, and p38.

Apoptosis Analysis^[3]

Cell Line:	OCI-AML3 cells
Concentration:	1 μ M
Incubation Time:	24 hours
Result:	Increased the levels of both cleaved PARP and cleaved caspase 3 and diminished bcl-2 and MCL1.

Cell Cycle Analysis^[3]

Cell Line:	OCI-AML3 cells
Concentration:	1 μ M
Incubation Time:	24 hours
Result:	Induced of G0-G1 arrest.

In Vivo

GNF-7 (10-20 mg/kg; o.p.; daily; for 7 days) displays significant in vivo efficacy against T315I Bcr-Abl in the bioluminescent xenograft mouse model^[2].

GNF-7 exhibits moderate oral bioavailability (mice 36%) and C_{max} (mice 3616 nM) following oral administration (mice 20 mg/kg)^[2].

GNF-7 exhibits terminal elimination half-lives (mice 3.8 h) due to high plasma clearance (8.6 mL/min/kg) following intravenous injection (mice 5 mg/kg)^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	6-8 weeks old SCID beige female mice, with Ba/F3-T315I-Bcr-Abl cells xenograft ^[2]
Dosage:	10 mg/kg, 20 mg/kg
Administration:	Oral administration, daily, for 7 days
Result:	Effectively inhibited tumor growth of T315I-Bcr-Abl-Ba/F3 cells in mice at low doses (10 mg/kg).

Animal Model:	5-6 weeks old male Balb/c mice (20-25 g) ^[2]
Dosage:	5 mg/kg for i.v.; 20 mg/kg for i.g. (Pharmacokinetic Analysis)
Administration:	Intravenous injection and oral gavage

Result:

Oral bioavailability (36%), C_{max} (3616 nM), T_{1/2} (3.2 h).

CUSTOMER VALIDATION

- Biochem Pharmacol. 2020 Jul;177:113947.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Choi HG, et al. A type-II kinase inhibitor capable of inhibiting the T315I "gatekeeper" mutant of Bcr-Abl. J Med Chem. 2010 Aug 12;53(15):5439-48.
- [2]. Lu X, et al. Hybrid pyrimidine alkynyls inhibit the clinically resistance related Bcr-Abl(T315I) mutant. Bioorg Med Chem Lett. 2015 Sep 1;25(17):3458-63.
- [3]. Cho, H., et al. First SAR study for overriding NRAS mutant driven acute myeloid leukemia. Journal of Medicinal Chemistry.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA