



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Alisertib

Cat. No.:	HY-10971
CAS No.:	1028486-01-2
Molecular Formula:	C ₂₇ H ₂₀ ClFN ₄ O ₄
Molecular Weight:	518.92
Target:	Aurora Kinase; Autophagy; Apoptosis
Pathway:	Cell Cycle/DNA Damage; Epigenetics; Autophagy; Apoptosis
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 1 year -20°C 6 months



SOLVENT & SOLUBILITY

In Vitro	DMSO : 25 mg/mL (48.18 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		1.9271 mL	9.6354 mL	19.2708 mL
		5 mM		0.3854 mL	1.9271 mL	3.8542 mL
		10 mM		0.1927 mL	0.9635 mL	1.9271 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.08 mg/mL (4.01 mM); Suspended solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.08 mg/mL (4.01 mM); Suspended solution; Need ultrasonic					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 2.08 mg/mL (4.01 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Alisertib (MLN 8237) is an orally active and selective Aurora A kinase inhibitor (IC ₅₀ =1.2 nM), which binds to Aurora A kinase resulting in mitotic spindle abnormalities, mitotic accumulation. Alisertib (MLN 8237) induces apoptosis and autophagy through targeting the AKT/mTOR/AMPK/p38 pathway in leukemic cells. Antitumor activity ^{[1][2][3]} .	
IC ₅₀ & Target	Aurora A 1.2 nM (IC ₅₀)	Aurora B 396.5 nM (IC ₅₀)

In Vitro	Alisertib (MLN 8237) leads the MM cells to mitotic spindle abnormalities, mitotic accumulation, as well as inhibition of cell proliferation through apoptosis and senescence. Alisertib up-regulates p53 and tumor suppressor genes p21 and p27^[1]. The decreased activity of Alisertib (MLN 8237) for the T217D/W277E Aurora A/TPX2 complex may reflect the increased affinity for ATP induced by cofactor binding to Aurora A^[2]. Alisertib (MLN 8237) inhibits cell proliferation with IC₅₀s ranging from 15 to 469 nM in different tumor cell lines^[4]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.								
In Vivo	Alisertib (MLN 8237) (30 mg/kg, p.o.) significantly reduces tumor burden and increases overall survival in xenograft-murine model of human-MM^[1]. Alisertib (3-30 mg/kg; p.o.; once daily for 3 weeks) causes tumor growth inhibition in solid tumor xenograft models^[4]. MCE has not independently confirmed the accuracy of these methods. They are for reference only. <table border="1"> <tr> <td>Animal Model:</td><td>Nude mice bearing HCT-116 colon tumor xenograft^[4]</td></tr> <tr> <td>Dosage:</td><td>3, 10, or 30 mg/kg</td></tr> <tr> <td>Administration:</td><td>P.o.; once daily for 3 weeks</td></tr> <tr> <td>Result:</td><td>Resulted in a dose-dependent TGI (tumor growth inhibition) of 43.3%, 84.2%, and 94.7% for the 3, 10, and 30 mg/kg groups, respectively.</td></tr> </table>	Animal Model:	Nude mice bearing HCT-116 colon tumor xenograft ^[4]	Dosage:	3, 10, or 30 mg/kg	Administration:	P.o.; once daily for 3 weeks	Result:	Resulted in a dose-dependent TGI (tumor growth inhibition) of 43.3%, 84.2%, and 94.7% for the 3, 10, and 30 mg/kg groups, respectively.
Animal Model:	Nude mice bearing HCT-116 colon tumor xenograft ^[4]								
Dosage:	3, 10, or 30 mg/kg								
Administration:	P.o.; once daily for 3 weeks								
Result:	Resulted in a dose-dependent TGI (tumor growth inhibition) of 43.3%, 84.2%, and 94.7% for the 3, 10, and 30 mg/kg groups, respectively.								

CUSTOMER VALIDATION

- Cell. 2017 Jan 12;168(1-2):264-279.e15.
- J Hepatol. 2021 Aug;75(2):363-376.
- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Theranostics. 2018 Feb 12;8(6):1740-1751.
- Clin Cancer Res. 2019 Jul 1;25(13):4179-4193.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Güllü G, et al. A novel Aurora-A kinase inhibitor MLN8237 induces cytotoxicity and cell-cycle arrest in multiple myeloma Blood June 24, 2010 vol. 115 no. 25 5202-5213.
- [2]. Sloane DA, et al. Drug-Resistant Aurora A Mutants for Cellular Target Validation of the Small Molecule Kinase Inhibitors MLN8054 and MLN8237 ACS Chem. Biol., 2010, 5 (6), pp 563-576.
- [3]. Bavetsias V, et al. Aurora Kinase Inhibitors: Current Status and Outlook. Front Oncol. 2015 Dec 21;5:278.
- [4]. Manfredi MG, et al. Characterization of Alisertib (MLN8237), an investigational small-molecule inhibitor of aurora A kinase using novel in vivo pharmacodynamic assays. Clin Cancer Res. 2011 Dec 15;17(24):7614-7624.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA