



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Tozadenant

Cat. No.:	HY-10995
CAS No.:	870070-55-6
Molecular Formula:	C ₁₉ H ₂₆ N ₄ O ₄ S
Molecular Weight:	406.5
Target:	Adenosine Receptor
Pathway:	GPCR/G Protein
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (123.00 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.4600 mL	12.3001 mL	24.6002 mL
		5 mM		0.4920 mL	2.4600 mL	4.9200 mL
		10 mM		0.2460 mL	1.2300 mL	2.4600 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.15 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Tozadenant is an adenosine A _{2A} receptor antagonist, with K _i of 11.5 nM on human A _{2A} and 6 nM on rhesus A _{2A} .
IC ₅₀ & Target	Ki: 6 nM (Rhesus A _{2A}), 11.5 nM (Human A _{2A})
In Vivo	¹⁸F-MNI-444 regional uptake is consistent with A_{2A} receptor distribution in the brain. Selectivity is demonstrated by dose-dependent blocking by tozadenant (1.5, 10.5 mg/kg) and preladenant^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Molecules. 2019 Apr 2;24(7):1295 .
- PLoS One. 2016 Nov 11;11(11):e0166415.
- PLoS One. 2016 Nov 11;11(11):e0166415.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Olivier Barret, et al. Adenosine 2A Receptor Occupancy by Tozadenant and Preladenant in Rhesus Monkeys. J Nucl Med. 2014 Oct;55(10):1712-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA