



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



GN44028

Cat. No.:	HY-110266
CAS No.:	1421448-26-1
Molecular Formula:	C ₁₈ H ₁₅ N ₃ O ₂
Molecular Weight:	305.33
Target:	HIF/HIF Prolyl-Hydroxylase
Pathway:	Metabolic Enzyme/Protease
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (163.76 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		3.2751 mL	16.3757 mL	32.7515 mL
		5 mM		0.6550 mL	3.2751 mL	6.5503 mL
		10 mM		0.3275 mL	1.6376 mL	3.2751 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (8.19 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (8.19 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (8.19 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	GN44028 is a potent and orally active hypoxia inducible factor (HIF)-1α inhibitor, with an IC ₅₀ of 14 nM. GN44028 inhibits hypoxia-induced HIF-1α transcriptional activity without suppressing HIF-1α mRNA expression, HIF-1α protein accumulation, or HIF-1α/HIF-1β heterodimerization. GN44028 can be used in the research of cancers ^{[1][3]} .
IC ₅₀ & Target	IC ₅₀ : 14 nM (HIF-1α) ^[1] .
In Vitro	GN44028 (compound 2l, 0-30 μM approximately) has anti-proliferative activities against HCT116, HepG2 and HeLa cells ^[1] .

GN44028 (0.001-1 μ M, 4 h) inhibits the hypoxia-induced VEGF mRNA expression in HeLa cells^[1].
 GN44028 (20 nM, 10 days) abrogates the TGF β induced colony formation in HCT116 cells^[2].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

RT-PCR^[1]

Cell Line:	HeLa cells
Concentration:	0.001, 0.01, 0.1, 1 μ M
Incubation Time:	4 h
Result:	Inhibited the hypoxia-induced HIF-1 transcriptional activity without affecting HIF-1 α /HIF-1 β heterodimerization.

In Vivo

GN44028 (10 mg/kg, oral gavage) extends the survival rate of animals in mice bearing a mixed orthotopic tumor with PN12 and ME23 cells^[3].
 GN44028 (5 mg/kg, tail vein injection, twice a week) suppresses tumor growth in a subcutaneous colorectal cancer model^[2].
 MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Mice bearing a mixed orthotopic tumor with PN12 and ME23 cells ^[3]
Dosage:	10 mg/kg
Administration:	Oral gavage
Result:	Reduced both proneural and mesenchymal markers.
Animal Model:	Subcutaneous colorectal cancer model (HCT116) ^[2]
Dosage:	5 mg/kg
Administration:	Tail vein injection, twice a week
Result:	Reduced pulmonary nodules in tumor-bearing mice, and enhanced the outcome of chemotherapy.

CUSTOMER VALIDATION

- ACS Nano. 2023 Nov 15.
- Oncogenesis. 2021 Oct 27;10(10):72.
- Environ Toxicol. 2021 May 10.
- J Pers Med. 2023, 13(1), 146.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Changfu Liu, et al. Tumor-associated macrophage-derived transforming growth factor- β promotes colorectal cancer progression through HIF1-TRIB3 signaling. Cancer Sci. 2021 Oct;112(10):4198-4207.
- [2]. Xiaoqing Fan, et al. Heterogeneity of subsets in glioblastoma mediated by Smad3 palmitoylation. Oncogenesis. 2021 Oct 27;10(10):72.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA