



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



VUF11207 fumarate

Cat. No.:	HY-110318
CAS No.:	1785665-61-3
Molecular Formula:	C ₃₁ H ₃₉ FN ₂ O ₈
Molecular Weight:	586.65
Target:	CXCR
Pathway:	GPCR/G Protein; Immunology/Inflammation
Storage:	4°C, sealed storage, away from moisture
	* In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (170.46 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	1.7046 mL	8.5230 mL	17.0459 mL
		5 mM	0.3409 mL	1.7046 mL	3.4092 mL
		10 mM	0.1705 mL	0.8523 mL	1.7046 mL
		Please refer to the solubility information to select the appropriate solvent.			
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution				
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution				
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (3.55 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	VUF11207 fumarate is a CXCR7 agonist that binds specifically to CXCR7. VUF11207 fumarate reduces CXCL12-mediated osteoclastogenesis and bone resorption by inhibiting ERK phosphorylation ^[1] .
IC₅₀ & Target	CXCR7 8.1 (pKi)
In Vitro	VUF11207 fumarate (0.17 nM; 5 days) inhibits RANKL- and TNF-α-induced osteoclastogenesis through CXCL12 inhibition in osteoclast precursor cells ^[1] . VUF11207 fumarate inhibits osteoclastogenesis by suppressing phosphorylation of erk ^[1] .

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[1]

Cell Line:	Osteoclast precursor cells (RANKL and Tnf α induced)
Concentration:	0.17 nM (100 ng/mL)
Incubation Time:	5 days
Result:	Showed inhibitory effect on CXCL12.

In Vivo

VUF11207 fumarate (100 μ g/day; s.c.; single daily for 5 days) inhibits LPS-induced osteoclastogenesis, bone resorption and production of RANKL and TNF α in mice^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Male c57Bl/6J wildtype/WT mice (8-10 week old; 20-25 g; LPS-induced) ^[1]
Dosage:	100 μ g/day
Administration:	Subcutaneous injection; single daily for 5 days
Result:	Significantly decreased the number of osteoclasts and suppressed Cathepsin K mRNA, ranKl and Tnf α mRNA expression levels. Reduced the area of LPS-induced bone resorption.

REFERENCES

[1]. Nugraha AP, et al. CXCR7 agonist acts as a CXCR7 motif chemokine ligand 12 inhibitor to ameliorate osteoclastogenesis and bone resorption. Mol Med Rep. 2022 Mar;25(3):78.

[2]. Wijnmans M, et al. Synthesis, modeling and functional activity of substituted styrene-amides as small-molecule CXCR7 agonists. Eur J Med Chem. 2012 May;51:184-92.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA