



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

AZ1495

Cat. No.:	HY-111101		
CAS No.:	2196204-23-4		
Molecular Formula:	C ₂₁ H ₃₁ N ₅ O ₂		
Molecular Weight:	385.5		
Target:	IRAK		
Pathway:	Immunology/Inflammation		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 10 mg/mL (25.94 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.5940 mL	12.9702 mL	25.9403 mL
		5 mM		0.5188 mL	2.5940 mL	5.1881 mL
		10 mM		0.2594 mL	1.2970 mL	2.5940 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1 mg/mL (2.59 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1 mg/mL (2.59 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil					
	Solubility: ≥ 1 mg/mL (2.59 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	AZ1495, a weak base, is a potent orally active interleukin-1 receptor associated kinase 4 (IRAK4) inhibitor. AZ1495 has a favorable physicochemical and kinase selectivity for IRAK4 and IRAK1 with IC ₅₀ values of 0.005 μM and 0.023 μM, respectively. AZ1495 has IRAK4 inhibition with a K _d value of 0.0007 μM. AZ1495 can be used for the research of diffuse large B-cell lymphoma (DLBCL) ^[1] .			
IC ₅₀ & Target	IRAK4 5 nM (IC ₅₀)	IRAK1 23 nM (IC ₅₀)	CLK1 50 nM (IC ₅₀)	CLK2 5 nM (IC ₅₀)

	CLK4 8 nM (IC ₅₀)	haspin 4 nM (IC ₅₀)
In Vitro	AZ1495 (compound 28) (10 µM, 1 h) has kinase selectivity for IRAK4 with IC₅₀ values of 0.005 µM (enzyme assay) and 0.052 µM (cellular assay), respectively^[1]. AZ1495 (10 µM, 1 h) has kinase inhibition for IRAK4 with an IC₅₀ value of 0.005 µM and K_d value of 0.0007 µM^[1]. AZ1495 (0.001-100 µM, 72 h) inhibits NF-κB activation and growth of ABC-DLBCL cell lines in a dosedependent manner^[1]. AZ1495 (0-3.3 µM, 14 h) completely inhibits NF-κB signaling and induces cell death at lower concentration in combination with a BTK inhibitor in OCI-LY10 cells^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only. Cell Viability Assay^[1]	
	Cell Line:	OCI-LY10 and SUDHL2 cells
	Concentration:	0.001-100 µM
	Incubation Time:	72 h
	Result:	Inhibited growth of OCI-LY10 cells in a dosedependent manner, whereas SUDHL2, a GCB-cell line was not sensitive and no increased cell killing to IRAK4 inhibitor. Increased the cell death in OCI-LY10 cells upon increasing concentrations of compound 28 and BTK ibrutinib.
	Western Blot Analysis ^[1]	
	Cell Line:	OCI-LY10 cells
	Concentration:	0-3.3 µM
	Incubation Time:	14 h
	Result:	Inhibited IκBα phosphorylation with dose-dependence in OCI-LY10 cells. Showed induction of apoptosis combination with 10 nM ibrutinib by cleavage of caspase 3 in OCI-LY10 cells.
In Vivo	AZ1495 (compound 28) (oral, daily, 12.5 mg/kg) leads to tumor regression combination with ibrutinib in an ABC-DLBCL mouse model (OCI-LY10 cells)^[1]. AZ1495 (iv., 2 mg/kg and oral, 5mg/kg) is characterized by high clearance (Cl) in rat (75 mL/min/kg) and moderate predictions based on hepatocyte data (Cl_{int} 15 µL/min/106 cells, predicted clearance 42 mL/min/kg) with low bioavailability consistent with a high first pass effect^[1]. AZ1495 (iv., 1 mg/kg) has low the amount of active renal secretion occurring in the dog^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	CB.17 SCID mice ^[1]
	Dosage:	12.5 mg/kg
	Administration:	oral, daily, 12.5 mg/kg
	Result:	Had modest anti-tumor activity as single agents but a combination of ibrutinib led to tumor regression and is well tolerated.
	Animal Model:	rat ^[1]

Dosage:	2 mg/kg, 5mg/kg																														
Administration:	iv., 2 mg/kg and oral, 5mg/kg																														
Result:	<table> <tr> <th>Species</th><th>Dose (mg/kg)</th><th>Cl (mL/min/kg)</th><th>Vss(L/kg)</th><th>PO halflife (h)</th><th>IV halflife (h)</th><th>Fabs (%)</th><th>F (%)</th></tr> <tr> <td>Rat</td><td>205</td><td>75</td><td>2.1</td><td>2.0</td><td>0.8</td><td>100</td><td>28</td></tr> <tr> <td>Dog</td><td>1</td><td>29</td><td>3.0</td><td>-</td><td>3.3</td><td>-</td><td>-</td></tr> </table>							Species	Dose (mg/kg)	Cl (mL/min/kg)	Vss(L/kg)	PO halflife (h)	IV halflife (h)	Fabs (%)	F (%)	Rat	205	75	2.1	2.0	0.8	100	28	Dog	1	29	3.0	-	3.3	-	-
Species	Dose (mg/kg)	Cl (mL/min/kg)	Vss(L/kg)	PO halflife (h)	IV halflife (h)	Fabs (%)	F (%)																								
Rat	205	75	2.1	2.0	0.8	100	28																								
Dog	1	29	3.0	-	3.3	-	-																								
Animal Model:	dog ^[1]																														
Dosage:	1 mg/kg																														
Administration:	iv., 1 mg/kg																														
Result:	<table> <tr> <th>Species</th><th>Dose (mg/kg)</th><th>Cl (mL/min/kg)</th><th>Vss(L/kg)</th><th>PO halflife (h)</th><th>IV halflife (h)</th><th>Fabs (%)</th><th>F (%)</th></tr> <tr> <td>Rat</td><td>205</td><td>75</td><td>2.1</td><td>2.0</td><td>0.8</td><td>100</td><td>28</td></tr> <tr> <td>Dog</td><td>1</td><td>29</td><td>3.0</td><td>-</td><td>3.3</td><td>-</td><td>-</td></tr> </table>							Species	Dose (mg/kg)	Cl (mL/min/kg)	Vss(L/kg)	PO halflife (h)	IV halflife (h)	Fabs (%)	F (%)	Rat	205	75	2.1	2.0	0.8	100	28	Dog	1	29	3.0	-	3.3	-	-
Species	Dose (mg/kg)	Cl (mL/min/kg)	Vss(L/kg)	PO halflife (h)	IV halflife (h)	Fabs (%)	F (%)																								
Rat	205	75	2.1	2.0	0.8	100	28																								
Dog	1	29	3.0	-	3.3	-	-																								

REFERENCES

[1]. Scott JS, et al. Discovery and Optimization of Pyrrolopyrimidine Inhibitors of Interleukin-1 Receptor Associated Kinase 4 (IRAK4) for the Treatment of Mutant MYD88L265P Diffuse Large B-Cell Lymphoma. J Med Chem. 2017 Dec 28;60(24):10071-10091.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA