



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HDAC8-IN-1

Cat. No.:	HY-111342
CAS No.:	1417997-93-3
Molecular Formula:	C ₂₂ H ₁₉ NO ₃
Molecular Weight:	345.39
Target:	HDAC
Pathway:	Cell Cycle/DNA Damage; Epigenetics
Storage:	Powder -20°C 3 years 4°C 2 years In solvent -80°C 2 years -20°C 1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (144.76 mM; ultrasonic and warming and heat to 60°C)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		2.8953 mL	14.4764 mL	28.9528 mL
		5 mM		0.5791 mL	2.8953 mL	5.7906 mL
		10 mM		0.2895 mL	1.4476 mL	2.8953 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.24 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (7.24 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (7.24 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	HDAC8-IN-1 is a HDAC8 inhibitor with an IC ₅₀ of 27.2 nM.
IC ₅₀ & Target	HDAC8 27.2 nM (IC ₅₀)
In Vitro	HDAC8-IN-1 is a HDAC8 inhibitor with an IC ₅₀ of 27.2 nM in cancer cell lines. HDAC8-IN-1 (compound 22 d) shows antiproliferative effects toward several human lung cancer cell lines (A549, H1299, and CL1-5); HDAC8-IN-1 exhibits

cytotoxicity against human lung CL1-5 cells without significant cytotoxicity for normal IMR-90 cells^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Death Dis. 2021 May 18;12(6):501.
- Metab Eng. 2023 Sep 17:80:94-106.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Huang WJ, et al. Synthesis and biological evaluation of ortho-aryl N-hydroxycinnamides as potent histone deacetylase (HDAC) 8 isoform-selective inhibitors. ChemMedChem. 2012 Oct;7(10):1815-24.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA