



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

BI-3812

Cat. No.:	HY-111381		
CAS No.:	2166387-64-8		
Molecular Formula:	C ₂₆ H ₃₂ ClN ₇ O ₅		
Molecular Weight:	558.03		
Target:	Bcl-2 Family		
Pathway:	Apoptosis		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 20.83 mg/mL (37.33 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg
		1 mM		1.7920 mL	8.9601 mL	17.9202 mL
		5 mM		0.3584 mL	1.7920 mL	3.5840 mL
		10 mM		0.1792 mL	0.8960 mL	1.7920 mL
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.08 mg/mL (3.73 mM); Suspended solution; Need ultrasonic					
	2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (3.73 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	BI-3812 is potent and efficacious BCL6 inhibitor, inhibiting the BTB domain of BCL6, with an IC ₅₀ of ≤3 nM; BI-3812 has antitumor activity.
IC ₅₀ & Target	IC ₅₀ : ≤3 nM (BCL6 BTB) ^[1]
In Vivo	BI-3812 is a BCL6 inhibitor, with an IC ₅₀ of ≤3 nM. BI-3812 shows an IC ₅₀ of 40 nM for the cellular BCL6 ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Kerres N, et al. Chemically Induced Degradation of the Oncogenic Transcription Factor BCL6. Cell Rep. 2017 Sep 19;20(12):2860-2875.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA